

Food insecurity and mental health outcomes among people living with HIV in West Africa

Running head: Food insecurity and mental health among PLHIV

Noelle A. BENZEKRI^{1*}, Marie K. PLAISY², Jean Jacques KOFFI³, Ephrem MENSAH⁵, Albert MINGA^{3,6}, Didier K. EKOUEVI^{2,4}, Raoul MOH³, Charlotte BERNARD², Antoine JAQUET² on behalf of the IeDEA West Africa Collaboration

¹ Departments of Medicine, Infectious Diseases, and Public Health, The University of Washington, Seattle WA, USA

² University of Bordeaux, National Institute for Health and Medical Research (INSERM) UMR 1219, Research Institute for Sustainable Development (IRD) EMR 271, Bordeaux Population Health Centre, Bordeaux, France

³ PAC-CI Research Program, Abidjan, Côte d'Ivoire

⁴ Faculté des Sciences de la Santé, Département de Santé Publique, Université de Lomé, Lomé, Togo

⁵ Espoir Vie Togo, Lomé, Togo

⁶ Centre Médical de Suivi de Donneurs de Sang (CMSDS), Abidjan, Cote d'Ivoire

* Corresponding author: benzekri@uw.edu

Disclaimers: The authors have no competing interests to declare.

Funding: The International Epidemiology Databases to Evaluate AIDS (IeDEA) West Africa is supported by grant number U01AI069919 from the United States National Institutes of Health (National Institute of Allergy and Infectious Diseases, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Heart, Lung, and Blood Institute, National Institute of Diabetes and Digestive and Kidney Diseases, National Institute on Drug Abuse, National Institute on Alcohol Abuse and Alcoholism, Fogarty International Center, and National Cancer Institute).

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Abstract

Objectives: This study was conducted in order to determine whether food insecurity is associated with poor mental health outcomes among PLHIV in Togo and Côte d'Ivoire, West Africa.

Design: Cross-sectional study among participants enrolled in the International epidemiological Databases to Evaluate AIDS West Africa cohort.

Methods: The Household Food Insecurity Access Scale was used to measure food insecurity. The PHQ-9, GAD-7, PCL-5, and AUDIT-C instruments were used to assess depressive symptoms, anxiety symptoms, symptoms of post-traumatic stress disorder (PTSD), and harmful alcohol use, respectively. Logistic regression was used to identify factors associated with poor mental health outcomes.

Results: Among 584 participants (68.8% female, median age 52 years), nearly half (48.1%) were food insecure; 7.0% were mildly food insecure, 26.7% were moderately food insecure, and 14.4% were severely food insecure. Severe food insecurity [OR 2.63 (1.30-5.34)] and being widowed [OR 2.15 (1.07-4.30)] were associated with moderate-to-severe depressive symptoms and formal education was protective [OR 0.30 (0.15-0.61)]; there was a trend towards an association between severe food insecurity and moderate-to-severe anxiety symptoms [OR 2.14 (0.98-4.70)] and formal education was protective [OR 0.35 (0.18-0.71)]; severe food insecurity [OR 6.87 (2.89-16.38)] and being widowed [OR 3.55 (1.46-8.59)] were associated with PTSD; mild food insecurity [OR 2.74 (1.20-6.26)] and male sex [OR 11.58 (5.44-24.68)] were associated with high risk alcohol use.

Conclusions: Food insecurity is associated with poor mental health outcomes among PLHIV in Togo and Côte d'Ivoire, West Africa. Future studies to understand and address the causal links between food insecurity and mental health among PLHIV are warranted.

Keywords: food insecurity; mental health; social determinants of health; HIV; West Africa; Sub-Saharan Africa; low- or middle-income countries

INTRODUCTION

According to the United Nations Food and Agriculture Organization, nearly a third of the global population is moderately or severely food insecure and does not have regular access to adequate food[1]. The major drivers of food insecurity include conflict, climate extremes, economic downturns, socio-structural inequality, and lack of access to nutritious foods and healthy environments[1]. These factors are particularly important in West Africa, where the prevalence of food insecurity continues to increase, and more than 60% of the population suffers from moderate or severe food insecurity[1].

The factors driving the increasing burden of food insecurity can also contribute to poor mental health. Globally, the prevalence of mental health disorders has increased by 48% in the past 30 years, with an estimated lifetime prevalence of any mental disorder of nearly 30%[2, 3]. According to the World Health Organization, the determinants of mental health include “not only individual attributes and interactions with others, but also social, cultural, economic, political and environmental factors”[4]. A

better understanding of the factors contributing to the increasing burden of mental health disorders is essential in order to both improve individual health outcomes and reverse this alarming trend.

Individuals living with HIV are uniquely impacted by both food insecurity and poor mental health[5-25]. Food insecurity is highly prevalent among people living with HIV (PLHIV) in West Africa and contributes to poor HIV outcomes, including poor adherence to antiretroviral therapy (ART), loss to follow-up, and virologic failure[17, 18, 26]. Due to a multitude of social, clinical, and biological factors, including HIV-associated stigma, advanced HIV disease, ART adverse effects, and socioeconomic stressors, PLHIV are also at increased risk of poor mental health[7, 8, 13, 27]. Prior studies suggest that depression is highly prevalent among PLHIV in West Africa[8], and may be as much as three times more prevalent among PLHIV compared to the general population[13, 28, 29]. Furthermore, mental health disorders such as depression, anxiety, and alcohol use disorder are associated with poor HIV outcomes, including increased risk of non-adherence to ART, loss to follow-up, virologic failure, and disease progression[6, 7, 30]. While food insecurity has been associated with poor mental health among members of the general population in West Africa, studies to specifically evaluate the relationship between food insecurity and mental health outcomes among PLHIV in the region are limited[31-40].

Understanding the associations between food insecurity and mental health among PLHIV may have important implications for the development and implementation of effective interventions to improve health outcomes and quality of life. This study was conducted in order to determine whether food insecurity is associated with poor mental health outcomes among PLHIV in Togo and Côte d'Ivoire, West Africa. The secondary objective was to determine how social determinants of health modify this effect.

METHODS

We conducted a cross-sectional study among a subset of participants enrolled in the International epidemiological Databases to Evaluate AIDS (IeDEA) West Africa cohort at the “Espoir Vie Togo (EVT)” Clinic in Lome, Togo and the “Centre Medical de Suivi des Donneurs de Sang (CMSDS)” in Abidjan, Côte d'Ivoire from November 2021 through June 2022. PLHIV who were ≥ 40 years old and receiving antiretroviral therapy for ≥ 6 months were enrolled as part of a larger study assessing the burden of non-communicable diseases in individuals aging with HIV across a variety of low-and-middle income settings[41, 42]. Participants were recruited during their routine visit using a systematic random sampling approach. Briefly, during a pre-enrollment visit and for a given day of HIV care visits, clinicians at the HIV clinic proposed participation to eligible patients using the following procedure: the research assistant in charge of enrollment selected every n th eligible patient (depending on the daily number of eligible attendees receiving HIV care), after randomly selecting the first patient through n th statistical elements as the starting point[43, 44]. Data were collected using structured questionnaires, validated instruments, and chart review.

Social determinants of health: Formal education, household income, and marital status were captured using a structured questionnaire.

Food Insecurity: The Household Food Insecurity Access Scale (HFIAS) was used to measure food insecurity[45]. Food secure was defined as HFIAS category 1 and food insecure was defined as HFIAS categories 2-4, where category 2 is mildly food insecure, category 3 is moderately food insecure, and category 4 is severely food insecure.

HIV outcomes: CD4 cell count and HIV viral load were captured using chart review.

Mental Health Outcomes: The Patient Health Questionnaire-9 (PHQ-9) instrument was used to assess severity of depressive symptoms, where a PHQ-9 score of 5-9 was categorized as mild depressive symptoms and a PHQ-9 score ≥ 10 was categorized as moderate-to-severe depressive symptoms[46-48]. The General Anxiety Disorder-7 (GAD-7) instrument was used to assess severity of anxiety symptoms, where a GAD-7 score of 5-9 was categorized as mild anxiety symptoms and a GAD-7 score ≥ 10 was categorized as moderate-to-severe anxiety symptoms[48, 49]. The PTSD Checklist for DSM-5 (PCL-5) instrument was used to assess symptoms of post-traumatic stress disorder (PTSD), where a PCL-5 score ≥ 33 was categorized as probable PTSD[50, 51]. The Alcohol Use Disorders Identification Test (AUDIT-C) instrument was used to assess harmful alcohol use, where a score ≥ 4 for men and ≥ 3 for women was categorized as harmful[52-54].

Statistical Analysis: Descriptive analysis was performed for all variables. Logistic regressions were used to identify predictors of mental health outcomes and to evaluate the associations between food insecurity and mental health outcomes using the categories that have been previously validated for each of these scales[46-54]. Variables that were significantly associated with mental health outcomes using simple regression were retained in the multivariable regression models. Missing data were excluded from analysis. P-values < 0.05 were considered significant. Data were analyzed using SPSS Statistics 29.

Ethical statement: All participants were informed about potential benefits and harms related to study participation and provided written informed consent prior to being included. The respective National Ethic committees of Cote d'Ivoire and Togo approved the study protocol (CMSDS, Côte d'Ivoire :195-21; EVT Clinic, Togo: 01/2022/CBRS).

RESULTS

CHARACTERISTICS OF STUDY PARTICIPANTS

Demographic characteristics: We enrolled 584 participants, of which 68.8% were female (**Table 1**). The median age was 52 years (IQR 48-58; range 41-80); 37.7% were 40-49 years of age, 54.3% were 50-64 years of age, and 8.0% were 65-80 years of age. Half of the participants were enrolled in Côte d'Ivoire and half were enrolled in Togo.

Social determinants of health: Nearly 17% had not received any formal education, 31.0% had attended primary school, 43.2% had attended secondary school, and 9.3% had attended university. Nearly 20% earned $< 25,000$ FCFA (< 42 USD)/month, 29.2% earned 25,000 to $< 50,000$ FCFA (42 to < 84 USD)/month, 23.5% earned 50,000 to $< 120,000$ FCFA (84 to < 202 USD)/month, and 27.7%

earned $\geq 120,000$ FCFA (≥ 202 USD)/month. More than half (51.2%) were married or cohabitating, 24.8% were single, 17.3% were widowed, and 6.7% were divorced or separated.

Food insecurity: Nearly half (48.1%) of participants were food insecure; 7.0% were mildly food insecure, 26.7% were moderately food insecure, and 14.4% were severely food insecure.

HIV outcomes: The median CD4 cell count was 583 cells/mm³ (IQR 396-816); only 6.6% of participants had a CD4 count < 200 cells/mm³. HIV viral loads were available for 562 participants, of which 90.7% had a viral load < 200 copies/mL.

Mental health outcomes: Moderate-to-severe depressive symptoms were experienced by 68 (11.7%) of participants, 44 (7.5%) of participants experienced moderate-to-severe anxiety symptoms, and 23 (3.9%) experienced PTSD. Harmful alcohol use was experienced by 132 (22.6%) of participants based upon the AUDIT-C score.

ASSOCIATIONS BETWEEN FOOD INSECURITY AND MENTAL HEALTH OUTCOMES

Depressive symptoms: Food insecurity (OR 2.11; 95% CI 1.24-3.59), particularly severe food insecurity (OR 3.35; 95% CI 1.77-6.31), was associated with moderate-to-severe depressive symptoms using simple logistic regressions (**Table 2**). Participants with an income $< 50,000$ FCFA (OR 1.89; 95% CI 1.08-3.30) and those who were widowed (OR 2.78; 95% CI 1.51-5.12) also had greater odds of moderate-to-severe depressive symptoms. Any level of formal education was protective (OR 0.27; 95% CI 0.15-0.49). Sex and age were not associated with moderate-to-severe depressive symptoms. In the multivariable regression model, severe food insecurity (OR 2.63; 95% CI 1.30-5.34) and being widowed (OR 2.15; 95% CI 1.07-4.30) were associated with moderate-to-severe depressive symptoms; formal education was protective (OR 0.30; 95% CI 0.15-0.61).

Anxiety symptoms: Severe food insecurity (OR 2.32; 95% CI 1.07-5.02) was associated with moderate-to-severe anxiety symptoms using simple logistic regressions and formal education was protective (OR 0.34; 95% CI 0.17-0.68) (**Table 3**). Sex, age, income, and marital status were not associated with moderate-to-severe anxiety symptoms. In the multivariable regression model, there was a trend towards an association between severe food insecurity and moderate-to-severe anxiety symptoms (OR 2.14; 95% CI 0.98-4.70); formal education was protective (OR 0.35; 95% CI 0.18-0.71).

PTSD: Food insecurity (OR 4.08; 95% CI 1.49-11.14), particularly severe food insecurity (OR 7.41; 95% CI 3.15-17.42), was associated with PTSD using simple logistic regressions (**Table 4**). Those who were widowed (OR 3.97; 95% CI 1.69-9.34)

also had greater odds of PTSD. Sex, age, education, and income were not associated with PTSD. In the multivariable regression model, severe food insecurity (OR 6.87; 95% CI 2.89-16.38) and being widowed (OR 3.55; 95% CI 1.46-8.59) were associated with PTSD.

Alcohol use: Based upon the AUDIT-C score, mild food insecurity (OR 2.45; 95% CI 1.24-4.82), male sex (OR 8.39; 95% CI 5.44-12.92), education beyond primary school (OR 1.57; 95% CI 1.06-

2.33), income \geq 50,000 FCFA (OR 1.83; 95% CI 1.21-2.76), and being married (OR 2.01; 95% CI 1.35-3.00) were associated with high risk alcohol use using simple regressions (**Table 5**). Age was not associated with high risk alcohol use. In the multivariable regression model, only mild food insecurity (OR 2.74; 95% CI 1.20-6.26) and male sex (OR 11.58; 95% CI 5.44-24.68) were associated with high risk alcohol use.

DISCUSSION

In this study conducted among PLHIV in Togo and Côte d'Ivoire, West Africa, we evaluated the impact of food insecurity on mental health outcomes. We found that food insecurity is common and is associated with depressive symptoms, anxiety symptoms, PTSD, and high risk alcohol use. To our knowledge, this is the first study to evaluate the impact of food insecurity on mental health outcomes among PLHIV in West Africa.

Individuals who are both food insecure and living with HIV are confronted by a multitude of biological and social factors that may heighten their risk of poor mental health outcomes. Food insecurity can contribute to poor mental health independent of HIV status as a result of nutrient deficiencies, increased levels of stress, feelings of shame, engaging in socially unacceptable coping strategies, disruptions to household ecology, loss of meal patterns and practices that sustain social cohesion, and increased burden of socioeconomic stressors [39, 55-57].

In the relationship between food insecurity and mental health, HIV can be a critical exacerbating factor through both direct and indirect pathways (**Figure 1**). HIV status can contribute to poor mental health outcomes due to factors such as HIV-associated stigma, clinical sequelae of advanced disease, side effects of ART, and HIV-associated socioeconomic stressors. These same HIV-associated factors can also lead to food insecurity or exacerbate preexisting food insecurity, thus further contributing to poor mental health through the food insecurity pathway. As such, the relationship between food insecurity and mental health may be amplified in the context of HIV.

Among the few studies that have evaluated the association between food insecurity and mental health among PLHIV in sub-Saharan Africa (SSA), the vast majority have focused on depression[58-66]. To our knowledge, only one previously published study has evaluated the association between food insecurity and anxiety symptoms among PLHIV in SSA, and there have been no prior studies that evaluated the association between food insecurity and PTSD in this context[64]. We were unable to identify any previously published studies conducted among PLHIV in SSA that used validated instruments to specifically evaluate the impact of food insecurity on high risk alcohol use[67, 68]. As such, this study helps to address a critical gap in the literature and makes an important contribution to our understanding of the complex relationship between food insecurity and mental health among PLHIV in SSA.

Depressive symptoms: Consistent with the findings from our study, food insecurity has been associated with depressive symptoms among PLHIV in other regions of SSA, including East Africa (Kenya[58, 59], Uganda[61, 62], Ethiopia[63], and Tanzania[64]) and Southern Africa (Zambia[65] and South Africa[66]). The majority of these studies used the HFIAS[58, 62, 65] or an adaptation of

the HFIAS to assess food insecurity[59, 63, 64]. There was greater variation in the instruments that were used to assess depression, which included the Hopkins Symptom Checklist for Depression (HSCL-D)[58, 62], the Center for Epidemiologic Studies Depression (CES-D) scale[59, 63, 66], the M.I.N.I. neuropsychiatric interview (MINI Plus) [61], the Children's Depression Inventory (CDI-S) [65], and the Edinburgh Postnatal Depression Scale (EPDS) [64]. In these studies food insecurity was associated with both greater odds of depression and greater severity of depressive symptoms.

Sociodemographic characteristics that were assessed included sex, age, marital status, education, occupation, household size, number of dependents, urban/rural residence, distance from residence to clinic, wealth, asset index, practice of agriculture, livestock ownership, ethnicity, religion, and social support. In a number of studies, women were more likely to experience depressive symptoms compared to men[58, 61-63], though this was not universal. Factors that were protective included marriage[64], livestock ownership[63], and financial capital[66].

Anxiety symptoms: In our study, there was a trend towards an association between food insecurity and anxiety symptoms. We were able to identify a single study conducted in SSA that evaluated the association between food insecurity and anxiety symptoms[64]. This study, which was conducted in Tanzania, used an adapted version of the HFIAS to assess food insecurity. Anxiety was assessed using the six-item subscale of the Brief Symptom Index (BSI-18), which differed from the instrument used in our study. Food insecurity was associated with greater odds of anxiety using simple regression, but not in the multivariable model where marriage was consistently protective.

Alcohol use: We found that food insecurity was associated with high risk alcohol use and men were more likely to engage in high risk alcohol use compared to women. To our knowledge, this is the first study conducted among PLHIV in SSA that used validated instruments to evaluate the association between food insecurity and high risk alcohol use. In a behavioral intervention study conducted among women living with HIV in South Africa who use alcohol, food insecurity was associated with less frequent alcohol use[67]. Importantly, neither food insecurity nor alcohol misuse were assessed using validated scales. It is of note that in our study, mild food insecurity but not moderate or severe food insecurity, was associated with high risk alcohol use. As food insecurity can be a correlate of poor economic status, it is possible that income is a factor which limits alcohol use among those experiencing severe food insecurity in some settings.

Social determinants: According to the WHO, the social determinants of health are “the non-medical factors that influence health outcomes”[69]. Among the social determinants evaluated in our study, the most important predictor of mental health outcomes was formal education. We found that formal education is protective against poor mental health outcomes, as individuals who had received a formal education were less likely to experience symptoms of depression or anxiety. An understanding of the social determinants that contribute to poor mental health outcomes, as well as those that are protective, is essential in order to develop and implement interventions that are informed by socio-cultural context.

Education is considered one of the most important modifiable social determinants of health[70]. It is strongly associated with life expectancy, health-seeking behaviors, and morbidity and is one of the most powerful tools by which individuals can emerge from poverty[70, 71]. However, in SSA it is

estimated that 105 million children of primary and secondary school age are not enrolled in school and only 23 percent of those enrolled complete secondary education[72]. Among the multitude of innovative and resource intensive health-associated interventions implemented in SSA, ensuring access to formal education should be universally prioritized.

Limitations: This study was conducted among older PLHIV on ART in urban centers in West Africa. As such, our findings may not be representative of all PLHIV or populations outside of this context. Our sample was limited to individuals ≥ 40 years of age. Greater age variability of our sample may have improved our ability to detect an association between increasing age and mental health outcomes. Participants in this study were PLHIV who were enrolled in care and receiving ART, which may have introduced selection bias. As both food insecurity and poor mental health are barriers to retention in care and adherence to ART, it is possible that our sample was biased towards those with greater food security and more favorable mental health outcomes. As such, the prevalence of food insecurity and mental health disorders in our study may be underestimated. While we were able to evaluate the associations between food insecurity and mental health outcomes, an understanding of the causal pathways was limited by the cross-sectional design. A future longitudinal study would help to elucidate the causal pathways between food insecurity and poor mental health and understand how the relationships between food insecurity and mental health change over time. The inclusion of additional social determinants of health would further strengthen future studies.

CONCLUSION

We found that food insecurity is associated with poor mental health outcomes among PLHIV in Togo and Côte d'Ivoire, West Africa. These findings suggest that interventions to address food insecurity may help to improve mental health outcomes for PLHIV. Future studies to understand and address the causal links between food insecurity and mental health are warranted.

ACKNOWLEDGEMENTS

Author contributions: Conceptualization: NB and AJ; Methodology: NB, AJ, and CB; Data collection: NB, MP, JJK; Writing – original draft: NB and AJ; Writing – revising and editing: NB, AJ, CB, MP, JJK; Resources and Supervision: DE, EM, AM, RM. All authors reviewed and approved the final manuscript.

The authors would like to thank the study participants and the staff of the Espoir Vie Togo (EVT) Clinic in Lome, Togo and the Centre Medical de Suivi des Donneurs de Sang (CMSDS) in Abidjan, Côte d'Ivoire.

The IeDEA West Africa Collaboration 2021-2026

Site investigators and cohorts

- Adult cohorts: Marcel Djimon Zannou, CNHU, Cotonou, Benin; Armel Poda, CHU Souro Sanou, Bobo Dioulasso, Burkina Faso; Oliver Ezechi, National Institute of Medical Research (NIMR), Lagos, Nigeria; Eugene Messou, ACONDA CePReF, Abidjan, Cote d'Ivoire; Henri

Chenal, CIRBA, Abidjan, Cote d'Ivoire; Kla Albert Minga, CMSDS, Abidjan, Cote d'Ivoire; Aristophane Tanon, CHU Treichville, Cote d'Ivoire; Moussa Seydi, CHU de Fann, Dakar, Senegal; Ephrem Mensah, Clinique EVT, Lomé, Togo.

- Pediatric cohorts: Caroline Yonaba, CHU Yalgado Ouadraogo, Ouagadougou, Burkina Faso; Lehila Bagnan Tossa, CNHU, Cotonou, Benin; Jocelyn Dame, Korle Bu Hospital, Accra, Ghana; Sylvie Marie N'Gbeche, ACONDA CePreF, Abidjan, Cote d'Ivoire; Kouadio Kouakou, CIRBA, Abidjan, Cote d'Ivoire; Madeleine Amorissani Folquet, CHU de Cocody, Abidjan, Cote d'Ivoire; François Tanoh Eboua, CHU de Yopougon, Abidjan, Cote d'Ivoire; Fatoumata Dicko Traore, Hopital Gabriel Toure, Bamako, Mali; Agatha David, NIMR, Lagos, Nigeria; Elom Takassi, CHU Sylvanus Olympio, Lomé, Togo.

Coordination & data centers

- Antoine Jaquet (PI), Didier Koumavi Ekouevi (PI), François Dabis, Renaud Becquet, Charlotte Bernard, Karen Malateste, Olivier Marcy, Marie Kerbie Plaisy, Elodie Rabourdin, Thierry Tiendrebeogo, ADERA, Isped & INSERM U1219/IRD, Bordeaux, France.
- Désiré Dahourou, Sophie Desmonde, Julie Jesson, Valeriane Leroy, Inserm U1295, Toulouse, France
- Raoul Moh, Jean-Claude Azani, Jean Jacques Koffi, Eric Komena, Maika Bengali, Abdoulaye Cissé, Guy Gnepa, Apollinaire Horo, Simon Boni, Eulalie Kangah, Corinne Moh, Jeanne Eliam, PACCI, CHU Treichville, Abidjan, Côte d'Ivoire

Partner institutions

- Ighovwerha Ofotokun (PI), Chris Martin, Emory University; Noelle Benzekri, Geoffrey Gottlieb, University of Washington; Olivia Keiser, Geneva University

Executive committee : Antoine Jaquet, Didier Ekouevi, Ighovwerha Ofotokun, François Dabis, Valérie Leroy

Steering committee : Renaud Becquet, Noelle Benzekri, Charlotte Bernard, Simon Boni, François Dabis, Désiré Dahourou, Sophie Desmonde, Didier Koumavi Ekouevi, Antoine Jaquet, Julie Jesson, Eric Komena, Valeriane Leroy, Karen Malateste, Ighovwerha Ofotokun, Marie Kerbie Plaisy, Elodie Rabourdin, Thierry Tiendrebeogo.

REFERENCES

- [1] Food and Agriculture Organization of the United Nations, UNICEF, WFP and WHO. 2024. The State of Food Security and Nutrition in the World 2024 – Financing to end hunger, food insecurity and malnutrition in all its forms. Rome.
- [2] "Global, regional, and national burden of 12 mental disorders in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019," (in eng), *Lancet Psychiatry*, vol. 9, no. 2, pp. 137-150, Feb 2022, doi: 10.1016/s2215-0366(21)00395-3.
- [3] J. J. McGrath *et al.*, "Age of onset and cumulative risk of mental disorders: a cross-national analysis of population surveys from 29 countries," (in eng), *Lancet Psychiatry*, vol. 10, no. 9, pp. 668-681, Sep 2023, doi: 10.1016/s2215-0366(23)00193-1.
- [4] "Comprehensive mental health action plan 2013–2030. Geneva: World Health Organization; 2021. ."
- [5] C. R. Rooks-Peck *et al.*, "Mental health and retention in HIV care: A systematic review and meta-analysis," (in eng), *Health Psychol*, vol. 37, no. 6, pp. 574-585, Jun 2018, doi: 10.1037/hea0000606.
- [6] J. Hoare, T. Sevenoaks, B. Mtukushe, T. Williams, S. Heany, and N. Phillips, "Global Systematic Review of Common Mental Health Disorders in Adults Living with HIV," (in eng), *Curr HIV/AIDS Rep*, vol. 18, no. 6, pp. 569-580, Dec 2021, doi: 10.1007/s11904-021-00583-w.
- [7] D. Chibanda, L. Benjamin, H. A. Weiss, and M. Abas, "Mental, Neurological, and Substance Use Disorders in People Living With HIV/AIDS in Low- and Middle-Income Countries," *JAIDS Journal of Acquired Immune Deficiency Syndromes*, vol. 67, pp. S54-S67, 2014, doi: 10.1097/qai.0000000000000258.
- [8] C. Bernard, F. Dabis, and N. de Rekeneire, "Prevalence and factors associated with depression in people living with HIV in sub-Saharan Africa: A systematic review and meta-analysis," *PLoS One*, vol. 12, no. 8, p. e0181960, 2017, doi: 10.1371/journal.pone.0181960.
- [9] P. N. Mwangala, A. Mabrouk, R. Wagner, C. Newton, and A. A. Abubakar, "Mental health and well-being of older adults living with HIV in sub-Saharan Africa: a systematic review," (in eng), *BMJ Open*, vol. 11, no. 9, p. e052810, Sep 22 2021, doi: 10.1136/bmjopen-2021-052810.
- [10] J. Hou *et al.*, "Posttraumatic Stress Disorder and Nonadherence to Treatment in People Living With HIV: A Systematic Review and Meta-analysis," (in eng), *Front Psychiatry*, vol. 11, p. 834, 2020, doi: 10.3389/fpsy.2020.00834.

- [11] F. H. Hu *et al.*, "Prevalence of mental health problems in people living with HIV: a systematic review and meta-analysis," (in eng), *Psychol Health Med*, pp. 1-17, Nov 6 2024, doi: 10.1080/13548506.2024.2424998.
- [12] L. C. Zalla *et al.*, "The anxiety care continuum and its association with viral suppression among persons with HIV," (in eng), *Aids*, vol. 38, no. 14, pp. 1956-1964, Nov 15 2024, doi: 10.1097/qad.0000000000003986.
- [13] S. M. Lofgren, D. J. Bond, N. Nakasujja, and D. R. Boulware, "Burden of Depression in Outpatient HIV-Infected adults in Sub-Saharan Africa; Systematic Review and Meta-analysis," (in eng), *AIDS Behav*, vol. 24, no. 6, pp. 1752-1764, Jun 2020, doi: 10.1007/s10461-019-02706-2.
- [14] T. A. Zeleke, K. Alemu, T. A. Ayele, Z. A. Denu, L. Mwanri, and T. Azale, "Systematic review and meta-analysis on the effect of depression on ART adherence among women living with HIV," (in eng), *PLoS One*, vol. 19, no. 6, p. e0300106, 2024, doi: 10.1371/journal.pone.0300106.
- [15] Y. Yang *et al.*, "Global prevalence of depressive symptoms among people living with HIV/AIDS: a systematic review and meta-analysis of the past five years," (in eng), *AIDS Care*, vol. 36, no. 2, pp. 153-164, Feb 2024, doi: 10.1080/09540121.2023.2285733.
- [16] M. Necho, Y. Zenebe, C. Tiruneh, G. Ayano, and B. Yimam, "The Global Landscape of the Burden of Depressive Symptoms/Major Depression in Individuals Living With HIV/AIDs and Its Effect on Antiretroviral Medication Adherence: An Umbrella Review," (in eng), *Front Psychiatry*, vol. 13, p. 814360, 2022, doi: 10.3389/fpsyt.2022.814360.
- [17] N. A. Benzekri *et al.*, "Food insecurity predicts loss to follow-up among people living with HIV in Senegal, West Africa," *AIDS Care*, vol. 34, no. 7, pp. 878-886, Jul 2022, doi: 10.1080/09540121.2021.1894316.
- [18] N. A. Benzekri *et al.*, "The impact of food insecurity on HIV outcomes in Senegal, West Africa: a prospective longitudinal study," *BMC Public Health*, vol. 21, no. 1, p. 451, Mar 6 2021, doi: 10.1186/s12889-021-10444-1.
- [19] N. T. Nkinsi, S. R. Galagan, N. A. Benzekri, S. Govere, and P. K. Drain, "Food Insecurity at HIV Diagnosis Associated with Subsequent Viremia Amongst Adults Living with HIV in an Urban Township of South Africa," (in eng), *AIDS Behav*, vol. 27, no. 11, pp. 3687-3694, Nov 2023, doi: 10.1007/s10461-023-04085-1.
- [20] S. D. Weiser *et al.*, "Longitudinal assessment of associations between food insecurity, antiretroviral adherence and HIV treatment outcomes in rural Uganda," (in eng), *Aids*, vol. 28, no. 1, pp. 115-20, Jan 2 2014, doi: 10.1097/01.aids.0000433238.93986.35.

- [21] "Young S, Wheeler AC, McCoy SI, Weiser SD. A review of the role of food insecurity in adherence to care and treatment among adult and pediatric populations living with HIV and AIDS. *AIDS and behavior*. 2014;18 Suppl 5:S505-15."
- [22] "Singer AW, Weiser SD, McCoy SI. Does Food Insecurity Undermine Adherence to Antiretroviral Therapy? A Systematic Review. *AIDS and behavior*. 2014."
- [23] W. Aibibula, J. Cox, A. M. Hamelin, T. McLinden, M. B. Klein, and P. Brassard, "Association Between Food Insecurity and HIV Viral Suppression: A Systematic Review and Meta-Analysis," (in eng), *AIDS Behav*, vol. 21, no. 3, pp. 754-765, Mar 2017, doi: 10.1007/s10461-016-1605-5.
- [24] E. Chop *et al.*, "Food insecurity, sexual risk behavior, and adherence to antiretroviral therapy among women living with HIV: A systematic review," (in eng), *Health Care Women Int*, vol. 38, no. 9, pp. 927-944, Sep 2017, doi: 10.1080/07399332.2017.1337774.
- [25] M. L. Hirsh *et al.*, "Food Insecurity Is Associated With Low Tenofovir Diphosphate in Dried Blood Spots in South African Persons With HIV," (in eng), *Open Forum Infect Dis*, vol. 10, no. 7, p. ofad360, Jul 2023, doi: 10.1093/ofid/ofad360.
- [26] N. A. Benzekri *et al.*, "High Prevalence of Severe Food Insecurity and Malnutrition among HIV-Infected Adults in Senegal, West Africa," *PLoS One*, vol. 10, no. 11, p. e0141819, 2015, doi: 10.1371/journal.pone.0141819.
- [27] S. Rueda *et al.*, "Examining the associations between HIV-related stigma and health outcomes in people living with HIV/AIDS: a series of meta-analyses," (in eng), *BMJ Open*, vol. 6, no. 7, p. e011453, Jul 13 2016, doi: 10.1136/bmjopen-2016-011453.
- [28] I. T. Gbadamosi *et al.*, "Depression in Sub-Saharan Africa," (in eng), *IBRO Neurosci Rep*, vol. 12, pp. 309-322, Jun 2022, doi: 10.1016/j.ibneur.2022.03.005.
- [29] M. C. Greene *et al.*, "The epidemiology of psychiatric disorders in Africa: a scoping review," (in eng), *Lancet Psychiatry*, vol. 8, no. 8, pp. 717-731, Aug 2021, doi: 10.1016/s2215-0366(21)00009-2.
- [30] A. Jaquet *et al.*, "Alcohol use and non-adherence to antiretroviral therapy in HIV-infected patients in West Africa," (in eng), *Addiction*, vol. 105, no. 8, pp. 1416-21, Aug 2010, doi: 10.1111/j.1360-0443.2010.02978.x.
- [31] J. P. Trudell, M. L. Burnet, B. R. Ziegler, and I. Luginaah, "The impact of food insecurity on mental health in Africa: A systematic review," (in eng), *Soc Sci Med*, vol. 278, p. 113953, Jun 2021, doi: 10.1016/j.socscimed.2021.113953.

- [32] K. N. Atuoye and I. Luginaah, "Food as a social determinant of mental health among household heads in the Upper West Region of Ghana," (in eng), *Soc Sci Med*, vol. 180, pp. 170-180, May 2017, doi: 10.1016/j.socscimed.2017.03.016.
- [33] G. Duthé, C. Rossier, D. Bonnet, A. B. Soura, and J. Corker, "Mental health and urban living in sub-Saharan Africa: major depressive episodes among the urban poor in Ouagadougou, Burkina Faso," (in eng), *Popul Health Metr*, vol. 14, p. 18, 2016, doi: 10.1186/s12963-016-0084-2.
- [34] R. M. Gyasi, B. Obeng, and J. Y. Yeboah, "Impact of food insecurity with hunger on mental distress among community-dwelling older adults," (in eng), *PLoS One*, vol. 15, no. 3, p. e0229840, 2020, doi: 10.1371/journal.pone.0229840.
- [35] A. Nyundo *et al.*, "Factors associated with depressive symptoms and suicidal ideation and behaviours amongst sub-Saharan African adolescents aged 10-19 years: cross-sectional study," (in eng), *Trop Med Int Health*, vol. 25, no. 1, pp. 54-69, Jan 2020, doi: 10.1111/tmi.13336.
- [36] M. Peele and S. Wolf, "Predictors of anxiety and depressive symptoms among teachers in Ghana: Evidence from a randomized controlled trial," (in eng), *Soc Sci Med*, vol. 253, p. 112957, May 2020, doi: 10.1016/j.socscimed.2020.112957.
- [37] A. C. Sweetland *et al.*, "Food insecurity, mental distress and suicidal ideation in rural Africa: Evidence from Nigeria, Uganda and Ghana," (in eng), *Int J Soc Psychiatry*, vol. 65, no. 1, pp. 20-27, Feb 2019, doi: 10.1177/0020764018814274.
- [38] C. Osei-Owusu, S. Dhillon, and I. Luginaah, "The impact of food insecurity on mental health among older adults residing in low- and middle-income countries: A systematic review," (in eng), *PLoS One*, vol. 19, no. 3, p. e0301046, 2024, doi: 10.1371/journal.pone.0301046.
- [39] L. Smith *et al.*, "Association between food insecurity and depression among older adults from low- and middle-income countries," (in eng), *Depress Anxiety*, vol. 38, no. 4, pp. 439-446, Apr 2021, doi: 10.1002/da.23147.
- [40] L. Smith *et al.*, "Association of food insecurity with suicidal ideation and suicide attempts in adults aged ≥ 50 years from low- and middle-income countries," (in eng), *J Affect Disord*, vol. 309, pp. 446-452, Jul 15 2022, doi: 10.1016/j.jad.2022.04.109.
- [41] A. M. Parcesepe *et al.*, "Prevalence and co-occurrence of symptoms of mental and substance use disorders among people with HIV age 40 and older in low- and middle-income countries: a cross-sectional study," (in eng), *J Int AIDS Soc*, vol. 27, no. 10, p. e26359, Oct 2024, doi: 10.1002/jia2.26359.

- [42] M. K. Plaisy *et al.*, "Metabolic causes of liver disease among adults living with HIV from low- and middle-income countries: a cross-sectional study," (in eng), *J Int AIDS Soc*, vol. 27, no. 4, p. e26238, Apr 2024, doi: 10.1002/jia2.26238.
- [43] R. J. Rossi, *Applied Biostatistics for the Health Sciences*. Newark, UNITED STATES: John Wiley & Sons, Incorporated, 2022.
- [44] M. H. Kuniholm *et al.*, "Association of cardiovascular disease risk with liver steatosis and fibrosis in people with HIV in low- and middle-income countries," *AIDS*, vol. 39, no. 1, pp. 11-21, Jan 1 2025, doi: 10.1097/QAD.0000000000004012.
- [45] "Coates J, Swindale A, Bilinsky P. Household Food Insecurity Access Scale (HFIAS) for Measurement of Household Food Access: Indicator Guide (v. 3). Washington, D.C.: Food and Nutrition Technical Assistance Project, Academy for Educational Development. 2007."
- [46] K. Kroenke, R. L. Spitzer, and J. B. Williams, "The PHQ-9: validity of a brief depression severity measure," (in eng), *J Gen Intern Med*, vol. 16, no. 9, pp. 606-13, Sep 2001, doi: 10.1046/j.1525-1497.2001.016009606.x.
- [47] D. Akena, J. Joska, E. A. Obuku, and D. J. Stein, "Sensitivity and specificity of clinician administered screening instruments in detecting depression among HIV-positive individuals in Uganda," *AIDS Care*, vol. 25, no. 10, pp. 1245-1252, 2013/10/01 2013, doi: 10.1080/09540121.2013.764385.
- [48] D. Chibanda *et al.*, "Validation of screening tools for depression and anxiety disorders in a primary care population with high HIV prevalence in Zimbabwe," (in eng), *J Affect Disord*, vol. 198, pp. 50-5, Jul 1 2016, doi: 10.1016/j.jad.2016.03.006.
- [49] R. L. Spitzer, K. Kroenke, J. B. Williams, and B. Lowe, "A brief measure for assessing generalized anxiety disorder: the GAD-7," *Arch Intern Med*, vol. 166, no. 10, pp. 1092-7, May 22 2006, doi: 10.1001/archinte.166.10.1092.
- [50] C. A. Blevins, F. W. Weathers, M. T. Davis, T. K. Witte, and J. L. Domino, "The Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5): Development and Initial Psychometric Evaluation," (in eng), *J Trauma Stress*, vol. 28, no. 6, pp. 489-98, Dec 2015, doi: 10.1002/jts.22059.
- [51] R. Verhey, D. Chibanda, L. Gibson, J. Brakarsh, and S. Seedat, "Validation of the posttraumatic stress disorder checklist - 5 (PCL-5) in a primary care population with high HIV prevalence in Zimbabwe," (in eng), *BMC Psychiatry*, vol. 18, no. 1, p. 109, Apr 23 2018, doi: 10.1186/s12888-018-1688-9.
- [52] K. Bush, D. R. Kivlahan, M. B. McDonell, S. D. Fihn, and K. A. Bradley, "The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol Use Disorders

Identification Test," (in eng), *Arch Intern Med*, vol. 158, no. 16, pp. 1789-95, Sep 14 1998, doi: 10.1001/archinte.158.16.1789.

- [53] K. A. Bradley, A. F. DeBenedetti, R. J. Volk, E. C. Williams, D. Frank, and D. R. Kivlahan, "AUDIT-C as a brief screen for alcohol misuse in primary care," (in eng), *Alcohol Clin Exp Res*, vol. 31, no. 7, pp. 1208-17, Jul 2007, doi: 10.1111/j.1530-0277.2007.00403.x.
- [54] S. Inoue *et al.*, "Testing the validity of the AUDIT-C and AUDIT-3 to detect unhealthy alcohol use among high-risk populations in Zambia: A secondary analysis from two randomized trials," (in eng), *Drug Alcohol Depend*, vol. 229, no. Pt A, p. 109156, Dec 1 2021, doi: 10.1016/j.drugalcdep.2021.109156.
- [55] L. J. Weaver and C. Hadley, "Moving beyond hunger and nutrition: a systematic review of the evidence linking food insecurity and mental health in developing countries," (in eng), *Ecol Food Nutr*, vol. 48, no. 4, pp. 263-84, Jul-Aug 2009, doi: 10.1080/03670240903001167.
- [56] A. D. Jones, "Food Insecurity and Mental Health Status: A Global Analysis of 149 Countries," *American Journal of Preventive Medicine*, vol. 53, no. 2, pp. 264-273, 2017, doi: 10.1016/j.amepre.2017.04.008.
- [57] S. Nanama and E. A. Frongillo, "Altered social cohesion and adverse psychological experiences with chronic food insecurity in the non-market economy and complex households of Burkina Faso," *Social Science & Medicine*, vol. 74, no. 3, pp. 444-451, 2012/02/01/ 2012, doi: 10.1016/j.socscimed.2011.11.009.
- [58] J. D. Miller *et al.*, "Household Water and Food Insecurity Are Positively Associated with Poor Mental and Physical Health among Adults Living with HIV in Western Kenya," (in eng), *J Nutr*, vol. 151, no. 6, pp. 1656-1664, Jun 1 2021, doi: 10.1093/jn/nxab030.
- [59] E. L. Tuthill *et al.*, "Persistent Food Insecurity, but not HIV, is Associated with Depressive Symptoms Among Perinatal Women in Kenya: A Longitudinal Perspective," (in eng), *AIDS Behav*, vol. 25, no. 3, pp. 847-855, Mar 2021, doi: 10.1007/s10461-020-03047-1.
- [60] G. O. Boateng, C. L. Workman, J. D. Miller, M. Onono, T. B. Neilands, and S. L. Young, "The syndemic effects of food insecurity, water insecurity, and HIV on depressive symptomatology among Kenyan women," (in eng), *Soc Sci Med*, vol. 295, p. 113043, Feb 2022, doi: 10.1016/j.socscimed.2020.113043.
- [61] E. Kinyanda, S. Hoskins, J. Nakku, S. Nawaz, and V. Patel, "Prevalence and risk factors of major depressive disorder in HIV/AIDS as seen in semi-urban Entebbe district, Uganda," (in eng), *BMC Psychiatry*, vol. 11, p. 205, Dec 30 2011, doi: 10.1186/1471-244x-11-205.
- [62] A. C. Tsai *et al.*, "Food insecurity, depression and the modifying role of social support among people living with HIV/AIDS in rural Uganda," (in eng), *Soc Sci Med*, vol. 74, no. 12, pp. 2012-9, Jun 2012, doi: 10.1016/j.socscimed.2012.02.033.

- [63] T. Yeneabat, A. Bedaso, and T. Amare, "Factors associated with depressive symptoms in people living with HIV attending antiretroviral clinic at Fitcha Zonal Hospital, Central Ethiopia: cross-sectional study conducted in 2012," (in eng), *Neuropsychiatr Dis Treat*, vol. 13, pp. 2125-2131, 2017, doi: 10.2147/ndt.S131722.
- [64] J. S. Ngocho *et al.*, "Depression and anxiety among pregnant women living with HIV in Kilimanjaro region, Tanzania," (in eng), *PLoS One*, vol. 14, no. 10, p. e0224515, 2019, doi: 10.1371/journal.pone.0224515.
- [65] S. Shangani, R. Masa, M. Zimba, G. Zimba, and D. Operario, "Food insecurity and depressive symptoms among young people living with HIV in Eastern Zambia," (in eng), *Int J STD AIDS*, vol. 35, no. 1, pp. 25-32, Jan 2024, doi: 10.1177/09564624231201917.
- [66] J. Hanass-Hancock, B. Carpenter, and H. Myezwa, "The missing link: exploring the intersection of gender, capabilities, and depressive symptoms in the context of chronic HIV," (in eng), *Women Health*, vol. 59, no. 10, pp. 1212-1226, Nov-Dec 2019, doi: 10.1080/03630242.2019.1607799.
- [67] F. A. Browne, M. W. Gichane, N. Shangase, J. Ndirangu, C. P. Bonner, and W. M. Wechsberg, "Social Determinants of Alcohol and Other Drug Misuse Among Women Living with HIV in Economically Underserved Communities in Cape Town, South Africa: A Cross-Sectional Study," (in eng), *AIDS Behav*, vol. 27, no. 4, pp. 1329-1338, Apr 2023, doi: 10.1007/s10461-022-03869-1.
- [68] A. M. Stanton *et al.*, "Factors Associated With Changes in Alcohol Use During Pregnancy and the Postpartum Transition Among People With HIV in South Africa and Uganda," (in eng), *J Int Assoc Provid AIDS Care*, vol. 22, p. 23259582231161029, Jan-Dec 2023, doi: 10.1177/23259582231161029.
- [69] World Health Organization. Health Topics: Social Determinants of Health. Available at: https://www.who.int/health-topics/social-determinants-of-health#tab=tab_1
- [70] H. The Lancet Public, "Education: a neglected social determinant of health," *The Lancet Public Health*, vol. 5, no. 7, p. e361, 2020, doi: 10.1016/S2468-2667(20)30144-4.
- [71] UNESCO, "The Right to Education." Available at: <https://www.unesco.org/en/right-education/need-know>
- [72] UNICEF, "Transforming Education in Africa: An evidence-based overview and recommendations for long-term improvements," 2021. [Online]. Available at: <https://www.unicef.org/media/106691/file/Transforming%20Education%20in%20Africa.pdf>

Figure 1. Relationships between food insecurity, mental health, and HIV status




Table 1. Characteristics of study participants, N=584

	N (%)
Female	402 (68.8)
Age, median years (IQR)[range]	52 (48-58)[41-80]
Age, years	
40-49	220 (37.7)
50-64	317 (54.3)
65-80	47 (8.0)
Country	
Côte d'Ivoire	292 (50)
Togo	292 (50)
Formal education	
None	96 (16.5)
Primary	181 (31.0)
Secondary	252 (43.2)
University	54 (9.3)

Income, FCFA/month*	
<25,000	103 (19.7)
25,000 to <50,000	153 (29.2)
50,000 to <120,000	123 (23.5)
≥120,000	145 (27.7)
Marital status	
Single	145 (24.8)
Married/cohabitating	299 (51.2)
Widowed	101 (17.3)
Divorced/separated	39 (6.7)
Food insecurity	
Food insecure	281 (48.1)
Mild food insecurity	41 (7.0)
Moderate food insecurity	156 (26.7)
Severe food insecurity	84 (14.4)
HIV outcomes	
Median CD4 cells/mm ³ (IQR)	583 (396-816)
CD4 <200 cells/mm ³	27 (6.6)
Viral suppression (<200 copies/mL) (<i>n</i> =562)	510 (90.7)
Depressive symptoms	
Mild depressive symptoms (PHQ-9 score 5-9)	178 (30.5)
Moderate-to-severe depressive symptoms (PHQ-9 score ≥10)	68 (11.7)
Anxiety symptoms	
Mild anxiety symptoms (GAD-7 score 5-9)	149 (25.5)
Moderate-to-severe anxiety symptoms (GAD-7 score ≥10)	44 (7.5)
Post-traumatic stress disorder (PTSD)	
Probable PTSD (PCL-5 score ≥33)	23 (3.9)
Harmful alcohol use	
Harmful alcohol use (AUDIT-C score ≥4 for men, ≥3 for women)	132 (22.6)

*25,000 FCFA = ~42 USD; 50,000 FCFA = ~84 USD; 120,000 FCFA = ~202 USD

Table 2. Logistic regressions: Predictors of moderate-to-severe depressive symptoms (PHQ-9 score ≥ 10)

	Univariable				Multivariable			
	OR	95% CI		p-value	OR	95% CI		p-value
Food Insecure (ref: not FI)	2.111	1.240	3.593	0.006				
Food Insecurity								
Mild FI	1.296	0.461	3.642	0.622				
Moderate FI	1.582	0.817	3.065	0.174				
Severe FI	3.941	1.980	7.844	<0.001				
Severe FI (ref: all others)	3.346	1.773	6.313	<0.001	2.630	1.296	5.335	0.007
Female	1.638	0.905	2.962	0.103				
Age (ref: 40-49)								
50-64	0.919	0.533	1.584	0.760				
65+	1.033	0.362	2.949	0.952				
Formal education (ref: none)								
Primary	0.319	0.155	0.653	0.002				
Secondary	0.286	0.146	0.561	<0.001				
University	0.079	0.017	0.356	<0.001				
Any formal education	0.267	0.145	0.491	<0.001	0.304	0.152	0.607	<0.001
Income (ref: 120,000+)								
<25,000	2.176	0.950	4.984	0.066				
25,000 to <50,000	2.019	0.958	4.252	0.065				
50,000 to <120,000	1.235	0.533	2.862	0.623				
Income <50,000	1.887	1.080	3.298	0.026	1.287	0.703	2.354	0.414
Marital status (ref: married)								
Single	1.475	0.770	2.822	0.241				
Widowed	3.165	1.628	6.154	<0.001				
Divorced/separated	1.044	0.291	3.752	0.947				
Widow (ref: all other)	2.775	1.507	5.108	<0.001	2.147	1.073	4.296	0.031

Table 3. Logistic regressions: Predictors of moderate-to-severe anxiety symptoms (GAD-7 score ≥ 10)

	Univariable				Multivariable			
	OR	95% CI		p-value	OR	95% CI		p-value
Food Insecure (ref: not FI)	1.148	0.616	2.142	0.664				
Food Insecurity								
Mild FI	0.559	0.126	2.485	0.445				
Moderate FI	0.913	0.417	2.000	0.821				
Severe FI	2.159	0.956	4.878	0.064				
Severe FI (ref: all others)	2.320	1.072	5.018	0.033	2.141	0.975	4.698	0.058
Female	1.019	0.522	1.989	0.956				
Age (ref: 40-49)								
50-64	0.658	0.346	1.248	0.200				
65+	0.376	0.085	1.674	0.199				
Formal education (ref: none)								
Primary	0.356	0.152	0.836	0.018				
Secondary	0.333	0.153	0.726	0.006				
University	0.299	0.080	1.113	0.072				
Any formal education	0.337	0.168	0.677	0.002	0.353	0.175	0.713	0.004
Income (ref: 120,000+)								
<25,000	0.432	0.151	1.236	0.118				
25,000 to <50,000	0.610	0.270	1.379	0.235				
50,000 to <120,000	0.410	0.153	1.095	0.075				
Income <50,000	0.752	0.382	1.480	0.410				
Marital status (ref: married)								
Single	1.380	0.647	2.943	0.404				
Widowed	1.218	0.513	2.890	0.655				
Divorced/separated	1.592	0.505	5.021	0.427				
Widow (ref: all other)	1.056	0.470	2.372	0.896				

Table 4. Logistic regressions: Predictors of PTSD (PCL-5 score 33+)

	Univariable				Multivariable			
	OR	95% CI		p-value	OR	95% CI		p-value
Food Insecure (ref: not FI)	4.079	1.494	11.139	0.006				
Food Insecurity								
Mild FI	0.000	0.000	-	0.998				
Moderate FI	2.384	0.716	7.938	0.157				
Severe FI	9.933	3.392	29.092	<0.001				
Severe FI (ref: all others)	7.409	3.152	17.417	<0.001	6.873	2.885	16.375	<0.001
Female	2.208	0.740	6.584	0.156				
Age (ref: 40-49)								
50-64	0.465	0.195	1.108	0.084				
65+	0.346	0.044	2.713	0.313				
Formal education (ref: none)								
Primary	2.006	0.546	7.370	0.294				
Secondary	1.148	0.304	4.334	0.838				
University	0.000	0.000	-	0.997				
Income (ref: 120,000+)								
<25,000	2.928	0.715	11.989	0.135				
25,000 to <50,000	1.932	0.474	7.873	0.358				
50,000 to <120,000	2.006	0.470	8.568	0.348				
Marital status (ref: married)								
Single	2.108	0.668	6.654	0.204				
Widowed	5.366	1.898	15.169	0.002				
Divorced/separated	1.285	0.151	10.964	0.819				
Widowed (ref: all other)	3.973	1.691	9.336	0.002	3.546	1.463	8.593	0.005

Table 5. Logistic regressions: Predictors of high risk alcohol use (AUDIT-C score ≥ 4 for men, ≥ 3 for women)

	Univariable				Multivariable			
	OR	95% CI		p-value	OR	95% CI		p-value
Food Insecure (ref: not FI)	1.019	0.691	1.502	0.923				
Food Insecurity								
Mild FI	2.448	1.243	4.819	0.010	2.736	1.196	6.262	0.017
Moderate FI	0.789	0.486	1.282	0.339				
Severe FI	0.943	0.524	1.695	0.843				
Male	8.385	5.444	12.915	<0.001	11.584	5.438	24.675	<0.001
Age, years (ref: 40-49)								
50-64	1.431	0.939	2.180	0.096				
65-80	1.145	0.528	2.487	0.731				
Formal education (ref: none)								
Primary	1.697	0.855	3.367	0.131				
Secondary	2.173	1.135	4.163	0.019				
University	2.688	1.176	6.143	0.019				
Education beyond primary school	1.569	1.055	2.332	0.026	0.805	0.431	1.503	0.496
Income (ref: <25,000)								
25,000 to <50,000	0.889	0.468	1.689	0.720				
50,000 to <120,000	2.000	1.079	3.707	0.028				
120,000+	1.474	0.799	2.720	0.215				
Income $\geq 50,000$	1.826	1.209	2.756	0.004	0.621	0.306	1.263	0.188
Marital status (ref: married)								
Single	0.525	0.319	0.864	0.011				
Widowed	0.510	0.286	0.909	0.022				
Divorced/separated	0.370	0.140	0.978	0.045				
Married (ref: all other)	2.011	1.347	3.004	<0.001	0.651	0.326	1.301	0.224