

RESEARCH

Open Access



Social determinants associated with psychological distress in children and adolescents during and after the first COVID-19-related lockdown in France: results from the CONFЕADO study

Mégane Estevez^{1*}, Nicolas Oppenchaim², Dalila Rezzoug^{3,4,5}, Isaura Laurent⁷, Sandrine Domecq¹, Imane Khireddine-Medouni⁸, Xavier Thierry⁹, Cédric Galera^{1,10}, Carla De Stefano^{3,4,6} and Stéphanie Vandentorren^{1,8,11*}

Abstract

Background This study aimed to analyze the parental socio-demographic characteristics of children and adolescents aged 9 to 18 years old, as well as the living and housing conditions associated with the psychological distress in these two sub-populations during and after France's first national COVID-19-related lockdown in spring 2020.

Methods We used data from the cross-sectional, observational, web-based study CONFЕADO, which collected data on children and adolescents' living and housing conditions and socio-demographic characteristics as well as those of their parents. It also collected data on children's and adolescents' health behaviors and psychological distress. We assessed psychological distress using the 10-item Children and Adolescents Psychological Distress Scale (CAPDS-10), and performed a multinomial logistic regression.

Results A total of 2882 children and adolescents were included in the present study. Factors associated with moderate psychological distress included being a female, parental financial difficulties, a lack of a private living space at home for the child/adolescent, and the following child health behaviors: no leisure or recreational activities with adults in the household, doing less than one hour of school homework a day, and not going outside during the lockdown. Severe psychological distress was associated with the parent's occupation (especially essential frontline workers), a lack of a private living space at home for the child/adolescent, and the following child health behaviors: spending over 5 h a day on social media, doing less than one hour of school homework a day, and no leisure or recreational activities with adults in the household.

Conclusions This study emphasizes the impact of housing and living conditions, as well as parents' socio-economic characteristics on children's health behaviors and psychological needs during the first COVID-19-related lockdown

*Correspondence:

Mégane Estevez

megane.estevez@gmail.com

Stéphanie Vandentorren

stephanie.vandentorren@santepubliquefrance.fr

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

in France. Our results suggest that health policies implemented during future pandemics should consider these structural social determinants to prevent severe psychological distress in children and adolescents.

Keywords COVID-19, Lockdown, Children, Adolescents, Living and housing conditions, Social determinants, Psychological distress

Background

In order to limit the spread of SARS-CoV-2 during the two years of the COVID-19 pandemic, most affected countries introduced unprecedented radical preventive measures, including lockdowns. This increased the burden of mental health in general populations, especially more isolated people, as well as those with difficult economic, housing, and family structure (e.g., single parents) conditions [1]. National strategies were also implemented to cope with the huge socioeconomic consequences of the pandemic (shorter working hours, increased unemployment benefit, etc.). In France, the government allowed shorter working hours at no cost to employers and implemented national lockdowns; the first lasted 8 weeks from 17 March to 11 May 2020. Schools were closed, with classes and homework continuing remotely. There were restrictions on family visits and local and international travel [2]. People had to remain at home; they were allowed outside for one hour a day provided they followed social distancing rules. Most socialization and care structures were closed [3].

This pandemic was a challenging period for children and adolescents (grouped under the single term ‘children’ where appropriate hereafter) worldwide, especially as late childhood and adolescence are critical periods for development and mental well-being [4]. School closures increased young people’s vulnerability to mental health problems, through reduced social interaction and physical activity, as well as increased screen time, irregular sleep patterns, and sub-optimal nutrition (especially in children for whom free school meals were an important source of nutrition [5], and in countries like France where the cost of fee-based school lunches was determined by family income [6]). These factors also negatively impacted children’s well-being in the short and long terms [7–11]. Younger people had few resources and few prior life experiences to be able to deal with stressful situations like COVID-19 [12]. A systematic review of the psychological impact of related lockdowns on children identified three studies reporting depression prevalence rates between 22.6% and 43.7%, and two studies reporting anxiety symptom rates between 18.9% and 37.4% [13].

Children from vulnerable social backgrounds, especially girls, were most affected by the impact of COVID-19 [14]. Parenting during the pandemic was an additional challenge for single families, for those on a low income, and those living in crowded housing. These families had to deal with remote learning (children), remote working

(parents), and a greater number of household chores [15]. Previous studies on the risk factors associated with poorer mental health among children and adolescents during the pandemic highlighted several predictors as follows: social isolation, screen time and excessive social media use, parental stress, a poor parent-child relationship, low socioeconomic status, preexisting mental health conditions, and/or preexisting disabilities [16]. However, in France, very few such studies have been conducted [17, 18], and none directly interviewed children or examined household-related structural social determinants (e.g., parent’s economic conditions and family structure) that could have affected their behavior and mental health during and after the pandemic.

We hypothesize that the living and housing conditions of many families, as well as the socio-demographic characteristics of parents during COVID-19-related lockdowns influenced children’s health behaviors, and had a considerable impact on the mental health of this population. Accordingly, we aimed to identify at-risk and protective structural social determinants associated with psychological distress in children and adolescents aged 9 to 18 years during and after France’s first COVID-19-related lockdown in spring 2020. The results from this study could inform both the development of interventions aimed at improving the mental health of children and adolescents’ during future pandemic-related lockdowns, and actions to address the issue of widening social inequity.

Methods

Design and study population

CONFÉADO (*CONFInement, Enfant et ADO*, translated as Lockdown, Child and Adolescent) is a cross-sectional observational study that examined how children experienced the first COVID-19-related lockdown in France. It was conducted between 9 June and 14 September 2020. Children from CONFÉADO who were in the care of child protection services were not included in the present analyses because data (parent-provided) on economic, living and housing conditions were not collected for this sub-population.

One of the child’s parents or his/her legal guardian had to provide consent for the child to participate. They also completed the parent questionnaire. Furthermore, children themselves had to provide their consent. In the present study, only children aged 9 to 18 years old who completed the 10-item Children and Adolescent’s

Psychological Distress Scale (CAPDS-10) for the assessment of psychological distress were included.

Data and procedure

CONFÉADO used a web-based anonymous, standardized questionnaire, which was completed by one of the participating child's parents (first part) and by the child him/herself (second part). The link to the questionnaire was sent to families by various institutions, associations, and partners of *University Sorbonne Paris Nord, Santé publique France* (SpF), *Observatoire national de la protection de l'enfance* (ONPE), *Assistance Publique-Hôpitaux de Paris* (AP-HP). It was also distributed on social networks.

Study variables

Children's psychological distress

The CAPDS-10 questionnaire was specifically designed for the CONFÉADO study and was validated in a previous study using data from CONFÉADO [19]. It is a self-assessment scale based on internationally validated scales including the Child Behavior Checklist (CBCL) and the Youth Self Report (YSR), and is intended to provide a measure of child psychological distress at a given time. All ten items concern a respondent's feelings over the previous two weeks. The first three assess his/her anxiety (nervousness/worry, stress, restlessness), the next three depression (lack of desire/pleasure in doing things, discouragement, lack of energy), the seventh somatic complaints (pain/fatigue/difficulty sleeping), and the remaining three aggressive behavior (opposition/irritability/disputes). Each item has four possible responses as follows: 'never', 'sometimes', 'more than half of the time', or 'almost every day', corresponding to a score between 0 (never) and 3 (almost every day). Accordingly, the total scale score ranges from 0 to 30; the higher the score, the more severe is the psychological distress. In the present study, the total score was analyzed as a three-modality categorical variable: scores between 0 and 9 indicated no or mild distress, scores between 10 and 18 indicated moderate distress, and scores between 19 and 30 indicated severe distress.

Socio-demographic characteristics of children and their parents

CONFÉADO collected data on two socio-demographic characteristics of the child's parents as follows: nationality (two French parents, one non-French parent, two non-French parents), and the level of education of the parent who completed the questionnaire (No diploma/Vocational training certificate, Upper secondary school certificate, Bachelor's degree, Master's degree/PhD/other).

For the present study, we constructed a categorical variable called 'parent's occupation' using data collected from an open-ended question about this theme in CONFÉADO which parents answered. In order to do this, we first created occupation clusters to group together occupations in the education sector (teachers, professors, etc.), healthcare professionals (doctors, nurses, psychologists, pharmacists, midwives, psychotherapists, etc.), essential frontline workers (housekeepers, mail carriers, drivers, cashiers, bakers, truck drivers, police officers, etc.), care providers (childcare providers, caregivers, nursing assistants, home aides, etc.), and commercial actors (florists, hairdressers, etc.). Occupations that did not fit into any of these six clusters and whose socio-professional category was known were added to one of the following three categories: 'other executive occupations', 'other intermediate occupations', 'other employee/manual worker occupations'.

The categorical variable 'parent's occupation' was then constructed as follows: 'frontline' and 'care providers' were grouped together into one modality (their commonality being that they continued to work after the lockdown started); 'healthcare professionals' and 'other occupations' comprised two extra modalities, with 'education sector' and 'commercial actors' included in the latter. Finally, there was an 'unknown' modality for those whose occupation was unknown.

CONFÉADO collected data on two socio-demographic characteristics of children: sex and age. Both were provided by the child (i.e., not by the parent) in the second part of the questionnaire (see above). Age was categorized into three age classes (9–11, 12–14, and 15–18 years) which correspond to different developmental stages of childhood and adolescence.

Living conditions

The two following variables were used: parent's self-perceived financial situation (very comfortable, comfortable, getting by, difficult to make ends meet) and family structure (two-parent, stepparent, single-parent). Both were provided by the parent (first part of questionnaire, see above).

Housing conditions

The three following variables were used: (i) municipal population density, categorized into 'highly', 'moderately', 'sparsely', and 'very sparsely' populated area; this variable was constructed from data on the child's place of residence, using the municipal density grid developed by the National Institute of Statistics and Economic Studies (INSEE; *Institut National de la Statistique et des Études Économiques*), which classifies districts in France into 4 categories based on municipal population density; (ii) type of housing (house with garden, house with

no garden/apartment/other); (iii) a private space in the accommodation if the child needed it (yes, no). The latter variable was provided by the child. The other two were provided by the parent.

Child mental health-related behaviors

Variables on the following child mental health-related behaviors during and after the lockdown were collected: how frequently the child went outside during the lockdown (never, approximately once a week, approximately 3 times a week, every day, once a day, multiple times a day), the time spent every day on social media since the beginning of the lockdown (none, less than an hour, 1 to 3 hours, 3 to 5 hours, more than 5 hours), the frequency of leisure and recreational activities with adults in the household since the beginning of the lockdown (including board games, video games, sports, and manual activities such as gardening, cooking, sewing, etc.) (never, approximately once a week, approximately 3 times a week, every day), and the time spent on school homework every day since the beginning of the lockdown (less than one hour, one hour, 2 to 3 hours, more than 3 hours). All these variables were provided by the children.

Child mental health and other variables

The following two variables were collected in CONFEADO and analyzed in the present study as they can be considered confounding factors in the relationship between social factors and psychological distress: history of child mental health disorder (yes, no) and child experience of a painful and/or stressful event (e.g., parents' divorce, death of a loved one, or a romantic disappointment) in the previous 12 months (yes, no). They were provided by the parent and the child, respectively. Data on COVID-19-related infections and hospitalizations of a loved one were also collected (provided by the child).

Statistical analysis

We developed a DAG (directed acyclic graph) (Fig. 1) to represent the relationships between the explanatory variables of interest, known confounding factors, and child psychological distress, using an etiological explanatory approach. The DAG allowed us to select a subset of relevant variables to include in our final model, according to the conceptual framework of the social determinants of COVID-19 on children's health and wellbeing [14].¹

We first conducted univariable analyses for each explanatory factor to explore the relationship between each independent variable and the dependent variable, ignoring the potential effect of other explanatory

variables. Then, we performed a multivariable multinomial logistic regression model on the complete dataset to investigate all explanatory variables, including the social factors associated with psychological distress. We were interested in the associations between parental socio-demographic characteristics (parent's occupation), living and housing conditions (variables that usually impact psychological distress, and may have influenced child health behaviors during the lockdown), age, sex, and whether a relative had been infected with COVID-19 and/or hospitalized for it. We adjusted for the parent's education level, nationality, child history of mental disorders, whether the child had experienced a painful and/or stressful event in the previous 12 months, and the type of housing. Application conditions were verified before running the model, including collinearity between variables and sufficient sample size. After running the model, fit and performance were evaluated. All analyses were performed using R version 4.2. The significance level was set at 5%.

Results

Sample characteristics

The present study included a total of 2882 children (Fig. 2; Table 1). Of these, 2005 (69.6%) were females, 393 (13.6%) were between 9 and 11 years old, 469 (16.3%) between 12 and 14 years old, and 2020 (70.1%) between 15 and 18 years old. Concerning child mental health, 887 (30.8%) and 201 (7%) had moderate and severe psychological distress, respectively, while 639 (22.2%) had a history of mental disorders; 1228 (42.6%) had had a painful and/or stressful event in the previous 12 months. In terms of child health behaviors, 536 (18.6%) went outside multiple times a day during the lockdown, 594 (20.6%) had spent more than 5 hours a day on social media since the lockdown began, and 643 (22.3%) had participated in leisure and recreational activities with adults in their household every day since the lockdown started. With regard to children's housing conditions, 438 (15.3%) did not have a private space in their living accommodation when needed. In terms of school work at home, 826 (28.7%) had spent more than 3 hours a day on school homework since the lockdown started. With regard to the impact of the COVID-19 pandemic on relatives, 910 (31.6%) had had a relative infected and/or hospitalized due to COVID-19.

In terms of parents' socio-demographic characteristics (Table 2), 211 (7.3%) were not French nationals. Concerning the respondent parent, 696 (24.1%) had no educational diploma, 348 (12.1%) were essential frontline workers, and 252 (8.7%) were healthcare professionals. With regard to living conditions, 686 (23.8%) were single-parent families, and 319 (11.1%) found it difficult to make ends meet. With regard to housing conditions,

¹ We performed a multivariable multinomial logistic regression model in an earlier version of the article before the DAG was developed following a reviewer's suggestion (data not shown but available on request).




Fig. 1 Directed acyclic graph (DAG) representing causal relationships between explanatory variables and child and adolescent psychological distress, CONFADO study, France, 2020

679 (23.6%) lived in a very sparsely populated area, and 881 (30.6%) lived in a house without a garden or an apartment.

Factors associated with psychological distress in univariable analysis

The univariable results revealed significant associations between moderate and/or severe psychological distress and several explanatory factors (Table 3) such as sex, age, two French parents, parent’s level of education, parental inactivity, single-parent family status, parental financial difficulties, type of housing, availability of private space within the living accommodation, history of mental health disorders, experiencing painful and/or stressful events in the previous 12 months, never going outside during the lockdown, spending more than one hour per day on social media since the beginning of the lockdown,

never engaging in leisure or recreational activities with adults in the household since the beginning of the lockdown, and spending less than one hour per day on school homework since the beginning of the lockdown. Additionally, having a loved one infected or hospitalized due to COVID-19 was also associated with moderate and severe psychological distress.

Factors remained associated with psychological distress in multivariable analysis

Risk factors for moderate psychological distress

The risk factors most strongly associated with moderate psychological distress (*versus* low or no psychological distress, i.e., the reference category) in the participating children are presented in Table 4. Multinomial logistic regression highlighted the following factors: being a female (aOR, 1.96 ; 95% CI, 1.57 ; 2.46), parental financial




Fig. 2 Flow chart of study sample, CONFEADO study, France, 2020

difficulties (aOR, 1.38 ; 95% CI, 1.04 ; 1.83), never going outside during the lockdown (aOR, 1.65 ; 95% CI, 1.15 ; 2.37), only going outside once a week (aOR, 1.48 ; 95% CI, 1.08 ; 2.03), or every day, once a day (aOR, 1.42 ; 95% CI, 1.05 ; 1.92), never doing leisure or recreational activities with adults in the household since the lockdown started (aOR, 1.53 ; 95% CI, 1.14 ; 2.04), and doing less than one

hour a day of school homework since the beginning of the lockdown (aOR, 1.42 ; 95% CI, 1.08 ; 1.88).

Protective factors for moderate psychological distress

The only protective factor for moderate psychological distress in children highlighted by multinomial logistic regression was the possibility of a private space within

Table 1 Characteristics of child and adolescent study participants, CONFEADO study, France, 2020. (N = 2882)

Variable	N (%)
Socio-demographic characteristics of children and adolescents	
Sex	2882
Male	877 (30.4)
Female	2005 (69.6)
Age (years)	2882
9–11	393 (13.6)
12–14	469 (16.3)
15–18	2020 (70.1)
Child and adolescent mental health	
Psychological distress	2882
No or mild distress	1794 (62.2)
Moderate distress	887 (30.8)
Severe distress	201 (7.0)
History of mental health disorder	2882
No	2243 (77.8)
Yes	639 (22.2)
Experience of a painful and/or stressful event in the previous 12 months	2882
No	1654 (57.4)
Yes	1228 (42.6)
Child mental health-related behaviors	
How frequently the child went outside during the first French lockdown	2882
Never	443 (15.4)
Approximately once a week	757 (26.3)
Approximately 3 times a week	514 (17.8)
Every day, once a day	632 (21.9)
Multiple times a day	536 (18.6)
Time spent every day on social media since the beginning of the first French lockdown	2882
None	394 (13.7)
Less than an hour	332 (11.5)
1 to 3 h	826 (28.7)
3 to 5 h	736 (25.5)
More than 5 h	594 (20.6)
Frequency of leisure and recreational activities with adults in the household since the beginning of the first French lockdown	2882
Never	648 (22.5)
Approximately once a week	862 (29.9)
Approximately 3 times a week	729 (25.3)
Every day	643 (22.3)
Time spent on school homework every day since the beginning of the first French lockdown	2882
Less than one hour	527 (18.3)
One hour	536 (18.6)
2 to 3 h	993 (34.5)
More than 3 h	826 (28.7)
Housing conditions	
A private space available in the accommodation if the child needed it	2882
No	438 (15.3)
Yes	2444 (84.8)
Impact of the COVID-19 pandemic on relatives	
COVID-19-related infection and/or hospitalization of a loved one	2882
No	1972 (68.4)
Yes	910 (31.6)

Table 2 Characteristics of the parents of children and adolescents included in the study sample, CONFEADO study, France, 2020. (N = 2882)

Variable	N (%)
Socio-demographic characteristics of parent	
<i>Parent's nationality</i>	
Two French parents	2882 2301 (79.8)
One non-French parent	370 (12.8)
Two non-French parents	211 (7.3)
<i>Parent's level of education</i>	
No diploma/Vocational training certificate	2882 696 (24.1)
Upper secondary school certificate	636 (22.1)
Bachelor's degree	735 (25.5)
Master's degree, PhD, other	815 (28.3)
<i>Parent's occupation</i>	
Essential frontline worker	2882 348 (12.1)
Healthcare professional	252 (8.7)
Other occupation	1681 (58.3)
Inactive	424 (14.7)
Unknown	177 (6.1)
Living conditions	
<i>Family structure</i>	
Two-parent	2882 1861 (64.6)
Stepparent	335 (11.6)
Single-parent	686 (23.8)
<i>Parent's financial difficulties</i>	
Very comfortable	2882 524 (18.2)
Comfortable	1100 (38.2)
Getting by	939 (32.6)
Difficult to make ends meet	319 (11.1)
Housing conditions	
<i>Municipal population density</i>	
High	2882 1086 (37.7)
Moderate	754 (26.2)
Sparse	363 (12.6)
Very sparse	679 (23.6)
<i>Type of housing</i>	
House with garden	2882 2001 (69.4)
House with no garden, apartment, other	881 (30.6)

the living accommodation when needed (aOR, 0.49; 95% CI, 0.38; 0.64).

Risk factors for severe psychological distress

The risk factors most strongly associated with severe psychological distress (Table 4) in children were: parent's occupation was a frontline worker (aOR, 1.66 ; 95% CI, 1.00 ; 2.75), spending more than 5 hours on social media a day since the lockdown started (aOR, 3.58 ; 95% CI, 1.38 ; 9.32), never doing any leisure and recreational

activity with adults in the household since the lockdown started (aOR, 2.71 ; 95% CI, 1.60 ; 4.59), and doing less than one hour a day of school homework since the beginning of the lockdown (aOR, 1.97 ; 95% CI, 1.24 ; 3.12).

Protective factors for severe distress

The only protective factor for severe psychological distress highlighted by the multinomial logistic regression was the possibility of a private space within the living accommodation when needed (aOR, 0.26; 95% CI, 0.17; 0.38).

Discussion

The results of the present study highlight the impact of structural social determinants, such as parents' socio-demographic characteristics, housing and living conditions, on children's psychological distress during the period of the first national lockdown. With regard to children's socio-demographic characteristics, being a female was associated with moderate psychological distress. Concerning parent socio-demographic characteristics, the parent's occupation - in particular being an essential frontline worker - was associated with severe psychological distress. In terms of living and housing conditions, parental financial difficulties were associated with moderate psychological distress. Furthermore, the possibility for the children to have a private space within the living accommodation when needed was a protective factor both for moderate and severe psychological distress. In terms of health behaviors, children with moderate distress in our study engaged less in physical activity (e.g., going outside); furthermore, the lack of leisure and recreational activities with adults in the household during the lockdown was a risk factor for both moderate and psychological distress. Doing less than one hour a day school homework since the beginning of the lockdown was associated with both moderate and severe psychological distress, while spending more than 5 hours a day on social media was associated with severe psychological distress.

In our study, as expected, being a female was associated with moderate psychological distress; this reflects findings from previous studies conducted in China [20], or in Brazil [21]. Even though females are usually more inclined to express their emotions, we hypothesize that the gendered distribution of domestic and educational tasks during the lockdown impacted females more, and that gender inequalities in terms of performing domestic tasks were accentuated [22, 23]. The national SAPRIS Elfe/Epipage2 survey, conducted in France in April-May 2020, found that girls were more involved in domestic chores in order to help parents cope with the increased pandemic-related domestic workload.

The results of our study show that children from socio-economically disadvantaged families were at a greater

Table 3 Factors associated with psychological distress in children and adolescents (9–18 years old) during and after the first COVID-19 pandemic-related lockdown in France in spring 2020: univariable multinomial logistic regression, CONFEADO study, France, 2020. (N = 3103)

Variable	Moderate distress vs. No or mild distress		Severe distress vs. No or mild distress	
	OR [95% CI]	p	OR [95% CI]	p
Children and adolescents socio-demographic characteristics				
<i>Sex</i>				
Male (Ref)	-		-	
Female	2.58 [2.14; 3.11]	< 0.001	2.48 [1.77; 3.47]	< 0.001
<i>Age (years)</i>				
9–11 (Ref)	-		-	
12–14	0.97 [0.70; 1.34]	0.853	1.89 [0.94; 3.81]	0.075
15–18	2.24 [1.74; 2.89]	< 0.001	4.19 [2.31; 7.61]	< 0.001
Parent's socio-demographic characteristics				
<i>Parent's nationality</i>				
Two French parents	0.80 [0.63; 1.00]	0.052	0.50 [0.35; 0.72]	< 0.001
One non-French parent (Ref)	-		-	
Two non-French parents	1.36 [0.96; 1.94]	0.088	1.37 [0.82; 2.29]	
<i>Parent's level of education</i>				
No diploma/Vocational training certificate (Ref)	-		-	
Upper secondary school certificate	0.75 [0.60; 0.94]	0.013	0.68 [0.47; 1.00]	0.049
Bachelor's degree	0.68 [0.55; 0.85]	< 0.001	0.55 [0.38; 0.81]	0.002
Master's degree, PhD, other	0.72 [0.58; 0.89]	0.002	0.61 [0.420; 0.87]	0.007
<i>Parent's occupation</i>				
Essential frontline worker	1.17 [0.92; 1.50]	0.208	1.98 [1.35; 2.91]	< 0.001
Healthcare professional	0.83 [0.63; 1.11]	0.221	0.94 [0.55; 1.59]	0.805
Inactive	1.29 [1.03; 1.60]	0.025	1.82 [1.27; 2.62]	0.001
Unknown	1.01 [0.73; 1.41]	0.942	1.25 [0.71; 2.22]	0.436
Other occupation (Ref)	-		-	
Living conditions				
<i>Family structure</i>				
Two-parent (Ref)	-		-	
Stepparent	1.06 [0.83; 1.37]	0.630	1.30 [0.84; 2.01]	0.239
Single-parent	1.23 [1.02; 1.48]	0.032	1.43 [1.03; 1.99]	0.032
<i>Parent's financial difficulties</i>				
Very comfortable (Ref)	-		-	
Comfortable	1.14 [0.90; 1.44]	0.270	1.59 [0.98; 2.58]	0.060
Getting by	1.83 [1.45; 2.30]	< 0.001	2.90 [1.81; 4.65]	< 0.001
Difficult to make ends meet	1.83 [1.36; 2.46]	< 0.001	3.88 [2.27; 6.63]	< 0.001
Housing conditions				
<i>Municipal population density</i>				
High	1.15 [0.95; 1.40]	0.157	1.28 [0.91; 1.81]	0.157
Moderate (Ref)	-		-	
Sparse	0.91 [0.69; 1.18]	0.473	0.59 [0.34; 1.04]	0.069
Very sparse	0.91 [0.73; 1.13]	0.395	1.16 [0.79; 1.70]	0.441
<i>Type of housing</i>				
House with garden	0.69 [0.58; 0.81]	< 0.001	0.59 [0.45; 0.79]	< 0.001
House with no garden, apartment, other (Ref)	-		-	
<i>A private space available in the accommodation if the child needed it</i>				
No (Ref)	-		-	
Yes	0.35 [0.29; 0.44]	< 0.001	0.16 [0.12; 0.21]	< 0.001
Child and adolescent mental health				
<i>History of mental health disorder</i>				
No (Ref)	-		-	
Yes	4.83 [3.9737; 5.87]	< 0.001	14.24 [10.52; 19.26]	< 0.001

Table 3 (continued)

Variable	Moderate distress vs. No or mild distress		Severe distress vs. No or mild distress	
	OR [95% CI]	p	OR [95% CI]	p
<i>Experience of a painful and/or stressful event in the previous 12 months</i>				
No (Ref)	-		-	
Yes	3.59 [3.05; 4.22]	< 0.001	8.55 [6.16; 11.86]	< 0.001
Child mental health-related behaviors				
<i>How frequently the child went outside during the lockdown</i>				
Never	2.98 [2.24; 3.97]	< 0.001	5.05 [2.99; 8.53]	< 0.001
Approximately once a week	2.49 [1.93; 3.22]	< 0.001	3.46 [2.09; 5.72]	< 0.001
Approximately 3 times a week	2.02 [1.53; 2.67]	< 0.001	1.72 [0.96; 3.09]	0.068
Every day, once a day	1.79 [1.36; 2.34]	< 0.001	2.13 [1.25; 3.64]	0.006
Multiple times a day (Ref)	-		-	
<i>Time spent every day on social media since the beginning of the lockdown</i>				
None (Ref)	-		-	
Less than an hour	1.04 [0.73; 1.47]	0.826	2.94 [1.33; 6.50]	0.008
1 to 3 h	1.54 [1.17; 2.04]	0.002	3.44 [1.68; 7.04]	< 0.001
3 to 5 h	2.24 [1.69; 2.96]	< 0.001	4.91 [2.41 10.03]	< 0.001
More than 5 h	3.24 [2.43; 4.33]	< 0.001	9.85 [4.86; 19.96]	< 0.001
<i>Frequency of leisure and recreational activities with adults in the household since the beginning of the lockdown</i>				
Never	2.24 [1.76; 2.84]	< 0.001	4.62 [3.06; 6.99]	< 0.001
Approximately once a week	1.55 [1.23; 1.94]	< 0.001	1.77 [1.15; 2.73]	0.010
Approximately 3 times a week	1.29 [1.02; 1.63]	0.032	0.70 [0.41; 1.20]	0.199
Every day (Ref)	-		-	
<i>Time spent on school homework every day since the beginning of the lockdown</i>				
Less than one hour	1.75 [1.39; 2.21]	< 0.001	3.02 [2.10 4.35]	< 0.001
One hour	1.08 [0.86; 1.36]	0.510	1.01 [0.66; 1.56]	0.960
2 to 3 h	1.01 [0.82; 1.23]	0.950	0.78 [0.53; 1.15]	0.210
More than 3 h (Ref)	-		-	
Impact of the COVID-19 pandemic on relatives				
<i>COVID-19-related infection and/or hospitalization of a loved one</i>				
No (Ref)	-		-	
Yes	1.30 [1.10; 1.54]	0.002	1.59 [1.20; 2.10]	0.001

OR: Odds Ratios; 95% CI: 95% Confidence Interval; p: p-value; Ref: Reference

risk of severe psychological distress. In univariable analysis, adolescents living with a single parent experienced more psychological distress than others (reflecting previous studies in France [25]). This factor was no longer significant in our multivariable model when financial difficulties were taken into account. The association between a parent being an essential frontline worker who had financial difficulties and psychological distress in the child confirms previous research indicating that financial hardship during the COVID-19 pandemic resulted in elevated stress levels while social inequalities strongly influenced the ways in which people experienced lockdowns [17, 23–33]. Similar associations between mental health and socio-economic factors during lockdowns have been found in other countries, such as in the UK [34] and in Canada [35].

Low-income households (which for the most part are immigrant households) living in poor housing conditions are overrepresented in the population of essential

frontline workers in France [36]. One of the French government's key measures during the pandemic was to prioritize the return to school of frontline healthcare workers' children, but not necessarily of other frontline workers. This may explain why unlike many studies which highlighted a negative influence of the COVID-19 pandemic on the mental health of frontline healthcare workers [37–40], having a parent who was a healthcare professional was not associated with children psychological distress in our multivariable model.

Our results regarding the association between the health behaviors of children and their psychological distress are consistent with those from a similar study in France [18]. That study found that living conditions were associated with a low health-related quality of life among children during the COVID-19 lockdown. Increased screen time became an integral element of adapting to the COVID-19 pandemic. A large study of children and

Table 4 Factors associated with psychological distress in children and adolescents (9–18 years old) during and after the first COVID-19 pandemic-related lockdown in France in spring 2020: multinomial logistic regression, CONFEADO study, France, 2020. (N = 2882)

Variable	Moderate distress (N = 887) vs. No or mild distress (N = 1794)		Severe distress (N = 201) vs. No or mild distress (N = 1794)	
	aOR [95% CI]	p	aOR [95% CI]	p
Children and adolescents socio-demographic characteristics				
<i>Sex</i>				
Male (Ref)	-		-	
Female	1.96 [1.57; 2.46]	< 0.001	1.52 [0.98; 2.37]	0.062
<i>Age (years)</i>				
9–11 (Ref)	-		-	
12–14	0.78 [0.53; 1.15]	0.205	1.19 [0.50; 2.80]	0.695
15–18	1.13 [0.79; 1.63]	0.505	0.90 [0.40; 2.01]	0.793
Parent's socio-demographic characteristics				
<i>Parent's occupation</i>				
Essential frontline worker	0.94 [0.69; 1.27]	0.693	1.66 [1.00; 2.75]	0.049
Healthcare professional	0.93 [0.66; 1.31]	0.691	0.96 [0.49; 1.87]	0.901
Inactive	0.91 [0.69; 1.21]	0.526	1.11 [0.67; 1.84]	0.677
Unknown	0.81 [0.55; 1.18]	0.272	1.25 [0.63; 2.48]	0.526
Other occupation (Ref)	-		-	
Living conditions				
<i>Family structure</i>				
Two-parent (Ref)	-		-	
Stepparent	0.75 [0.56; 1.00]	0.072	0.85 [0.51; 1.42]	0.536
Single-parent	0.81 [0.64; 1.02]	0.072	0.67 [0.44; 1.01]	0.056
<i>Parent's financial difficulties</i>				
Very comfortable (Ref)	-		-	
Comfortable	1.02 [0.78; 1.34]	0.882	1.23 [0.70; 2.16]	0.461
Getting by	1.38 [1.04; 1.83]	0.025	1.51 [0.86; 2.68]	0.153
Difficult to make ends meet	1.03 [0.70; 1.50]	0.896	1.08 [0.53; 2.18]	0.832
Housing conditions				
<i>Municipal population density</i>				
High	0.97 [0.76; 1.23]	0.785	1.01 [0.64; 1.58]	0.980
Moderate (Ref)	-		-	
Sparse	1.00 [0.74; 1.36]	0.997	0.86 [0.44; 1.66]	0.648
Very sparse	0.96 [0.73; 1.24]	0.734	1.51 [0.93; 2.44]	0.098
<i>A private space available in the accommodation if the child needed it</i>				
No (Ref)	-		-	
Yes	0.49 [0.38; 0.64]	< 0.001	0.26 [0.17; 0.38]	< 0.001
Child mental health-related behaviors				
<i>How frequently the child went outside during the lockdown</i>				
Never	1.65 [1.15; 2.37]	0.006	1.53 [0.78; 3.01]	0.215
Approximately once a week	1.48 [1.08; 2.03]	0.015	1.40 [0.75; 2.61]	0.293
Approximately 3 times a week	1.37 [0.99; 1.90]	0.057	0.86 [0.43; 1.71]	0.665
Every day, once a day	1.42 [1.05; 1.92]	0.025	1.27 [0.68; 2.38]	0.454
Multiple times a day (Ref)	-		-	
<i>Time spent every day on social media since the beginning of the lockdown</i>				
None (Ref)	-		-	
Less than an hour	0.71 [0.47; 1.07]	0.101	2.16 [0.81; 5.75]	0.124
1 to 3 h	1.01 [0.70; 1.47]	0.939	2.40 [0.94; 6.08]	0.066
3 to 5 h	1.17 [0.79; 1.72]	0.435	2.54 [0.98; 6.56]	0.055
More than 5 h	1.41 [0.94; 2.12]	0.098	3.58 [1.38; 9.32]	0.009
<i>Frequency of leisure and recreational activities with adults in the household since the beginning of the lockdown</i>				
Never	1.53 [1.14; 2.04]	0.004	2.71 [1.60; 4.59]	< 0.001
Approximately once a week	1.23 [0.95; 1.61]	0.119	1.40 [0.82; 2.37]	0.217

Table 4 (continued)

Variable	Moderate distress (N=887) vs. No or mild distress (N=1794)		Severe distress (N=201) vs. No or mild distress (N=1794)	
	aOR [95% CI]	p	aOR [95% CI]	p
Approximately 3 times a week	1.10 [0.84; 1.45]	0.476	0.61 [0.33; 1.13]	0.117
Every day (Ref)	-	-	-	-
<i>Time spent on school homework every day since the beginning of the lockdown</i>				
Less than one hour	1.42 [1.08; 1.88]	0.012	1.97 [1.24; 3.12]	0.004
One hour	1.10 [0.83; 1.44]	0.510	0.85 [0.50; 1.45]	0.555
2 to 3 h	1.12 [0.89; 1.42]	0.321	0.95 [0.60; 1.50]	0.827
More than 3 h (Ref)	-	-	-	-
Impact of the COVID-19 pandemic on relatives				
<i>COVID-19-related infection and/or hospitalization of a loved one</i>				
No (Ref)	-	-	-	-
Yes	1.04 [0.86; 1.26]	0.697	1.21 [0.85; 1.71]	0.286

aOR: adjusted Odds Ratios; 95% CI: 95% Confidence Interval; p: p-value; Ref: Reference

The results were adjusted for parent's nationality, parent's level of education, type of housing, history of mental health disorder and experience of a painful and/or stressful event in the previous 12 months

adolescents also linked smartphone addiction and excessive internet use to increased depression [41].

In our study, not having an appropriate place to self-isolate at home was related both to moderate and severe psychological distress. This result is congruent with that of Gatell-Carbó et al. who showed that children and adolescents with abnormal SDQ (Strengths and Difficulties Questionnaire) scores had substantially less personal space to study than the rest of their sample [43].

France's first lockdown significantly disrupted school routines. Some parents, especially single parents, had to juggle between the huge burden of childcare, household chores, and teleworking which may have negatively impacted the time they could dedicate to homeschooling. Studies have highlighted that children from low-income families were less likely to have access to digital tools and a good quality internet connection to do their homework; they were also less likely to have an isolated room to work in during the lockdown [42, 43]. Consequently, distance learning during the pandemic aggravated educational inequalities in France and elsewhere [44, 45].

Although infection and/or hospitalization of a relative due to COVID-19 was not a factor for psychological distress in our multivariable model, its impact should not be ignored, as many studies have shown that it can affect the mental health status of young people [31, 41, 46–48]. In univariable analysis, infection and/or hospitalization of a relative due to COVID-19 was associated both with moderate and severe psychological distress, but was no longer significant in our multivariable model when being an essential frontline worker was taken into account. Indeed, previous studies have shown that COVID-19 related morbidity and mortality was higher in migrants [49], overrepresented in frontline workers [36].

Strengths and limitations

The study has several limitations. First, a part of the data collection occurred during the two-month summer school holiday period (July-August 2020) in France when children would not normally have been at school. Ideally, data collection should have been concluded by the end of June. Second, school closure experiences were reported retrospectively; we had no data on when exactly the child returned to school or for how many hours per week. This made it impossible to directly measure the influence of returning to school on children mental health. Third, CONDEADO did not collect data on parents' mental health. These could have been useful, especially for mothers, as they may have a greater influence on the emotional well-being of their children. Stress and anxiety levels related to the health crisis may have affected children's mental health. Fourth, we chose to include children whose parents did not provide information about their occupation (the 'unknown' occupation category, see above). The presence of this 'unknown' modality in this variable may have led to potential bias in our results. Fifth, CONFEADO was conducted without a sample design and it is probable that families interested in mental health were more inclined to agree to participate. Nevertheless, considering a diverse range of social situations - through the collection of socio-demographic and environmental data from parents - allowed us to take into account social health inequalities.

Thus, our study has also several strengths. It is the first national study to assess the mental health and experiences of children during and after the first lockdown in France using self-evaluation. Furthermore, the CAPDS-10 scale has good psychometric characteristics and provides a reliable measure of distress at a given time [19]. Given its cross-sectional design, this study acknowledges

the limitations in drawing causal relationships, but it still offers valuable evidence and sheds light on potential links between social determinants such as socio-economic circumstances and children's psychological distress, during the pandemic contributing to our understanding of their mental health during this unprecedented crisis.

Conclusions

Our results highlight the importance of having socially differentiated public policies, in particular to support and accompany families in fragile socioeconomic conditions in order that the needs of all children can be met. This was especially true for low-income frontline workers during the COVID-19 pandemic who had to continue to work outside of home while everyone else was expected to stay at home. In the event of future lockdowns (for whatever reason), educational continuity and the earliest possible return to the classroom are two actions which could help limit feelings of being overwhelmed by school work, and the related negative impact on the mental health of children. Protective factors such as the quality of relationships and screen-free activities within the family need to be reinforced through accessible and appropriate information for parents and children.

List of Abbreviations

aOR	Adjusted Odds Ratio
AP-HP	Assistance Publique-Hôpitaux de Paris
ASE	Aide Sociale à l'Enfance
CAPDS-10	Children and Adolescent's Psychological Distress Scale
CBCL	Child Behavior Checklist
CI	Confidence Interval
CNIL	Commission nationale de l'informatique et des libertés
CONFÉADO	CONFInement, Enfant et ADO
INSEE	Institut National de la Statistique et des Études Économiques
ONPE	Observatoire national de la protection de l'enfance
SDQ	Strengths and Difficulties Questionnaire
SpF	Santé publique France
YSR	Youth Self Report

Acknowledgements

The authors thank the CONFÉADO consortium (Myriam Vimond, Emeline Delaville, Anne Oui, Sarah Gensburger, Agnès Gindt-Ducros, Carole Dufouil, Anne Lise Denoued, Nolwen Regnault) and Thierry Baubet, Fabien d'Hont and Bruno Falissard. Special thanks to Enguerrand Habran for participating in the recruitment of participants, overseeing the statistical analyses, and discussing the results. Our thanks to our colleagues in Santé publique France Aude Coivous, Anne-Laure Mereau, Suzanne Montanary and Sophie Lamarque for assisting with the dissemination of the study. Thanks also to Sahar Haidar for translation support and to Jude Sweeney (Milan, Italy) for copyediting the English version. Finally, our thanks to all the children and parents who participated in the study.

Authors' contributions

ME, SV, NO wrote the manuscript. ME, NO, DR, XT, CG, CDS, and SD contributed materials and analysis tools. ME, SV, NO, and SD analyzed the data. SV, NO, CDS, DR, and SD conceived and designed the study. SV and NO supervised the study. All authors contributed to the article, critically revised the manuscript for important intellectual content, and approved the submitted version.

Funding

None to declare.

Data Availability

The datasets generated and/or analyzed during the current study are not publicly available due to limitations associated with ethical approval. They are available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

An information note was provided to the children before they completed the questionnaire (with the objectives of the research, the anonymity and confidentiality of the data, the lack of obligation to answer the questions, etc.), as well as a parent consent form. The CONFÉADO study was approved by the CPP (Committee for the Protection of Persons n°2020-A01342.37). The study and all involved methods were carried out in accordance with relevant guidelines and regulations. The data collection and processing implemented in this research complies with the reference methodology MR003 approved by the CNIL (Commission nationale de l'informatique et des libertés).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no conflict of interest.

Author details

- ¹Centre de recherche Bordeaux Population Health (BPH), Université de Bordeaux, Inserm U1219, Bordeaux, France
- ²Université de Tours, UMR CITERES 7324, 35 Allée Ferdinand de Lesseps, Tours 37200, France
- ³Université Paris 13 Sorbonne Paris Nord, Bobigny, France
- ⁴AP-HP, Department of Child and Adolescent Psychiatry and General Psychiatry, Avicenne Hospital, Bobigny, France
- ⁵Centre de Recherche en épidémiologie et santé des populations (CESP), Université Paris Saclay, Villejuif, France
- ⁶AP-HP, Urgences - Samu 93, Hôpital Avicenne, Bobigny, France
- ⁷École nationale de la statistique et de l'analyse de l'information (ENSAI), Bruz, France
- ⁸Santé publique France, Saint Maurice, France
- ⁹ELFE Joint Unit, French Institute for Demographic Studies (Ined), French Institute for Medical Research and Health (Inserm), Paris, France
- ¹⁰Department of Child and Adolescent Psychiatry, Centre Hospitalier Charles Perrens, Bordeaux, France
- ¹¹Centre National de Ressources et de Résilience Lille-Paris (CN2R), Lille, France

Received: 27 September 2022 / Accepted: 10 July 2023

Published online: 18 July 2023

References

1. Fianu A, Aissaoui H, Naty N, et al. Health Impacts of the COVID-19 Lockdown measure in a low Socio-Economic setting: a cross-sectional study on Reunion Island. *Int J Environ Res Public Health*. 2022;19:13932.
2. Santé publique France. COVID-19 chez l'enfant (moins de 18 ans). État des lieux de la littérature en amont de la réouverture annoncée des crèches et des écoles. État de la littérature au 24 avril 2020. Synthèse rapide COVID-19, / import/covid-19-chez-l-enfant-moins-de-18-ans-.etat-des-lieux-de-la-litterature-en-amont-de-la-reouverture-annoncee-des-creches-et-des-ecoles-etat-de-l (accessed 9 March 2021).
3. Pinel-Jacquemin S, Martinez A, Martinasso M, et al. Lockdown in France: impact on families of Young Children with Special needs. *Front Psychol*. 2022;13:781030.
4. Santé mentale des adolescents., <https://www.who.int/fr/news-room/fact-sheets/detail/adolescent-mental-health> (accessed 20 July 2021).
5. Viner R, Russell S, Saull R, et al. School Closures during Social Lockdown and Mental Health, Health Behaviors, and well-being among children and adolescents during the First COVID-19 Wave: a systematic review. *JAMA Pediatr*. 2022;176:400–9.

6. Dubuisson C, Lioret S, Dufour A, et al. Socio-economic and demographic variations in school lunch participation of french children aged 3–17 years. *Public Health Nutr.* 2011;14:227–38.
7. Wang G, Zhang Y, Zhao J, et al. Mitigate the effects of home confinement on children during the COVID-19 outbreak. *Lancet Lond Engl.* 2020;395:945–7.
8. Ghosh R, Dubey MJ, Chatterjee S, et al. Impact of COVID – 19 on children: special focus on the psychosocial aspect. *Minerva Pediatr.* 2020;72:226–35.
9. Imran N, Zeshan M, Pervaiz Z. Mental health considerations for children & adolescents in COVID-19 pandemic. *Pak J Med Sci.* 2020;36:67–572.
10. Guessoum SB, Lachal J, Radjack R, et al. Adolescent psychiatric disorders during the COVID-19 pandemic and lockdown. *Psychiatry Res.* 2020;291:113264.
11. Li W, Wang Z, Wang G, et al. Socioeconomic inequality in child mental health during the COVID-19 pandemic: first evidence from China. *J Affect Disord.* 2021;287:8–14.
12. Mohler-Kuo M, Dzemaili S, Foster S, et al. Stress and Mental Health among Children/Adolescents, their parents, and young adults during the First COVID-19 Lockdown in Switzerland. *Int J Environ Res Public Health.* 2021;18:4668.
13. Nearchou F, Flinn C, Niland R, et al. Exploring the impact of COVID-19 on Mental Health Outcomes in Children and Adolescents: a systematic review. *Int J Environ Res Public Health.* 2020;17:E8479.
14. Lemkow-Tovías G, Lemkow L, Cash-Gibson L, et al. Impact of COVID-19 inequalities on children: an intersectional analysis. *Sociol Health Illn.* 2023;45:145–62.
15. de Figueiredo CS, Sandre PC, Portugal LCL, et al. COVID-19 pandemic impact on children and adolescents' mental health: Biological, environmental, and social factors. *Prog Neuropsychopharmacol Biol Psychiatry.* 2021;106:110171.
16. Caffo E, Asta L, Scandroglio F. Predictors of mental health worsening among children and adolescents during the coronavirus disease 2019 pandemic. *Curr Opin Psychiatry.* 2021;34:624–30.
17. Monnier M, Moulin F, Thierry X et al. Children's Mental and behavioral health, schooling, Sociodemographic and socioeconomic characteristics during School Closure in France due to COVID-19: the SAPRIS Project. Epub ahead of print 7 October 2021. DOI: <https://doi.org/10.21203/rs.3.rs-568392/v1>.
18. Bourion-Bédès S, Rousseau H, Batt M, et al. The effects of living and learning conditions on the health-related quality of life of children and adolescents during the COVID-19 lockdown in the French Grand Est region. *BMC Public Health.* 2022;22:517.
19. De Stefano C, Rezzoug D, Laurent I et al. Children and Adolescents Psychological Distress Scale during COVID-19 pandemic: validation of a psychometric instrument (CONFEADO study). *Front Psychiatry.*
20. Zhou S-J, Zhang L-G, Wang L-L, et al. Prevalence and socio-demographic correlates of psychological health problems in chinese adolescents during the outbreak of COVID-19. *Eur Child Adolesc Psychiatry.* 2020;29:749–58.
21. Garcia de Avila MA, Hamamoto Filho PT, Jacob FL da. S, Children's anxiety and factors related to the COVID-19 pandemic: an exploratory study using the children's anxiety questionnaire and the Numerical Rating Scale. *Int J Environ Res Public Health;* 17. Epub ahead of print 09 2020. DOI: <https://doi.org/10.3390/ijerph17165757>.
22. Alon T, Doepke M, Olmstead-Rumsey J et al. *The Impact of COVID-19 on Gender Equality.* crctr24_2020_163, University of Bonn and University of Mannheim, Germany, https://ideas.repec.org/p/bon/boncr/crctr24_2020_163.html (March 2020, accessed 24 March 2021).
23. Thierry X, Geay B, Pailhé A et al. Les enfants à l'épreuve du premier confinement. *Popul Soc* 2021; N° 585: 1–4.
24. Ezpeleta L, Navarro JB, de la Osa N et al. Life Conditions during COVID-19 Lockdown and Mental Health in Spanish Adolescents. *Int J Environ Res Public Health;* 17. Epub ahead of print 07 2020. DOI: <https://doi.org/10.3390/ijerph17197327>.
25. Recchi E, Ferragina E, Godechot O et al. Living through Lockdown: Social Inequalities and Transformations during the COVID-19 Crisis in France, https://spire.sciencespo.fr/hdl/2441/6vv2fug6nb8t29ilm995n9hbnh/resources/op-2020-1.pdf?_ga=2.182867046.1950713339.1604399123-1433383576.1604399123 (2020, accessed 24 March 2021).
26. Saurabh K, Ranjan S. Compliance and psychological impact of Quarantine in Children and Adolescents due to Covid-19 pandemic. *Indian J Pediatr.* 2020;87:532–6.
27. Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *The Lancet.* 2020;395:912–20.
28. Clemens V, Deschamps P, Fegert JM et al. Potential effects of “social” distancing measures and school lockdown on child and adolescent mental health. *Eur Child Adolesc Psychiatry* 2020; 1–4.
29. Cai Q, Feng H, Huang J, et al. The mental health of frontline and non-frontline medical workers during the coronavirus disease 2019 (COVID-19) outbreak in China: a case-control study. *J Affect Disord.* 2020;275:210–5.
30. Tso WYW, Wong RS, Tung KTS et al. Vulnerability and resilience in children during the COVID-19 pandemic. *Eur Child Adolesc Psychiatry.* Epub ahead of print 2020. DOI: <https://doi.org/10.1007/s00787-020-01680-8>.
31. Luijten MAJ, van Muilekom MM, Teela L, et al. The impact of lockdown during the COVID-19 pandemic on mental and social health of children and adolescents. *Qual Life Res Int J Qual Life Asp Treat Care Rehabil.* 2021;30:2795–804.
32. Sanzeri O, Zampini C, Belle R, et al. Premier confinement: 392 000 travailleurs de « première ligne », en majorité des femmes. INSEE; 2021. <https://www.insee.fr/fr/statistiques/5895960>.
33. Sugg MM, Runkle JD, Andersen L, et al. Crisis response among essential workers and their children during the COVID-19 pandemic. *Prev Med.* 2021;153:106852.
34. Mansfield KL, Newby D, Soneson E, et al. COVID-19 partial school closures and mental health problems: a cross-sectional survey of 11,000 adolescents to determine those most at risk. *JCPP Adv.* 2021;1:e12021.
35. Polsky JY, Gilmour H. Food insecurity and mental health during the COVID-19 pandemic. *Health Rep.* 2020;31:3–11.
36. Acs M, Chevrot J, Beaufrils S et al. *Quelles conditions de travail et de vie pour les 1,8 million de travailleurs « essentiels du quotidien » résidant en Île-de-France ?* INSEE, <https://www.insee.fr/fr/statistiques/5401737> (2021).
37. Feng Z, Xu L, Cheng P, et al. The psychological impact of COVID-19 on the families of first-line rescuers. *Indian J Psychiatry.* 2020;62:438–S444.
38. Gupta S, Sahoo S. Pandemic and mental health of the front-line healthcare workers: a review and implications in the indian context amidst COVID-19. *Gen Psychiatry.* 2020;33:e100284.
39. Spoorthy MS, Pratapa SK, Mahant S. Mental health problems faced by healthcare workers due to the COVID-19 pandemic—A review. *Asian J Psychiatry.* 2020;51:102119.
40. Billings J, Ching BCF, Gkofa V, et al. Experiences of frontline healthcare workers and their views about support during COVID-19 and previous pandemics: a systematic review and qualitative meta-synthesis. *BMC Health Serv Res.* 2021;21:923.
41. Duan L, Shao X, Wang Y, et al. An investigation of mental health status of children and adolescents in china during the outbreak of COVID-19. *J Affect Disord.* 2020;275:112–8.
42. Lambert A, Cayouette-Remblière J, Guérait É, et al. Le travail et ses aménagements: ce que la pandémie de covid-19 a changé pour les Français. *Popul Soc.* N° 2020;579:1–4.
43. Cullinane C, Montacute R. COVID-19 and Social mobility impact brief #1: School Shutdown. The Sutton Trust; 2020.
44. Delès R, Pirone F, Rayou P. L'accompagnement scolaire pendant le premier confinement de 2020. *Adm Educ* 2021; N° 169: 155–61.
45. Goudeau S, Sanrey C, Stanczak A et al. Why lockdown and distance learning during the COVID-19 pandemic are likely to increase the social class achievement gap. *Nat Hum Behav* 2021; 1–9.
46. Cao W, Fang Z, Hou G, et al. The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry Res.* 2020;287:112934.
47. Yeasmin S, Banik R, Hossain S, et al. Impact of COVID-19 pandemic on the mental health of children in Bangladesh: a cross-sectional study. *Child Youth Serv Rev.* 2020;117:105277.
48. Mangolian Shahrabaki P, Dehghan M, Maazallahi M et al. Fear and anxiety in girls aged 7 to 11 years old and related factors during the coronavirus pandemic. *Clin Child Psychol Psychiatry* 2021; 13591045211013872.
49. Khat M, Ghosn W, Guillot M, et al. Impact of the COVID-19 crisis on the mortality profiles of the foreign-born in France during the first pandemic wave. *Soc Sci Med.* 2022;313:115160.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.