






OPEN Parents' depression and anxiety associated with hyperactivity-inattention and emotional symptoms in children during school closure due to COVID-19 in France

Maëva Monnier^{1,2}, Flore Moulin¹, Marion Bailhache^{1,3}, Xavier Thierry^{4,5,6}, Stéphanie Vandentorren^{1,7}, Sylvana Côté^{1,2,8}, Bruno Falissard^{9,10}, Thierry Simeon^{4,5,6}, Bertrand Geay^{4,5,6}, Laetitia Marchand-Martin^{5,11,12,13}, Marie-Noelle Dufourg^{4,5,6}, Marie-Aline Charles^{4,5,6,11,12}, Pierre-Yves Ancel^{5,11,12,13}, Maria Melchior¹⁴, Alexandra Rouquette^{9,10}, Cédric Galera^{1,2,8}, The SAPRIS Study Group* & The ELFE Joint Unit*

Several risk factors of children's mental health issues have been identified during the pandemic of CoronaVirus Disease first appeared in 2019 (COVID-19). This study aims to fill the knowledge gap regarding the association between parents' and children's mental health issues during the COVID-19 school closure in France. We conducted a cross-sectional analysis of data collected in the SAPRIS-ELFE study during the COVID-19 pandemic in France. Using multinomial logistic regressions, we estimated associations between parents' and children's mental health issues. Symptoms of anxiety were assessed by the General Anxiety Disorder-7 (GAD-7) and depression by the Patient Health Questionnaire-9 (PHQ-9) for the parents. Hyperactivity/inattention and emotional symptoms in children were assessed by the Strengths and Difficulties Questionnaire (SDQ). The sample included 3496 children aged 8 to 9 years, of whom 50.0% were girls. During the school closure, 7.1% of responding parents had moderate to severe levels of anxiety and 6.7% had moderate to severe levels of depression. A total of 11.8% of the children had an abnormal hyperactivity/inattention score and 6.6% had an abnormal emotional symptoms score. In multivariate regression models, parental moderate to severe level of anxiety and moderate to severe level of depression were associated with abnormal hyperactivity-inattention score (adjusted Odds Ratio (aOR) 3.31; 95% Confidence Interval (CI) 2.33–4.70 and aOR 4.65; 95% CI 3.27–6.59, respectively) and abnormal emotional symptoms score in children (aOR 3.58; 95% CI 2.33–5.49 and aOR 3.78; 95% CI 2.47–5.78 respectively). Children whose

¹Bordeaux Population Health Research Center, Institut National de la Santé et de la Recherche Médicale (INSERM) U 1219, 146 Rue Léo Saignat, 33077 Bordeaux Cedex, France. ²Department of Child and Adolescent Psychiatry, Centre Hospitalier Charles Perrons, Bordeaux, France. ³Pole de Pédiatrie, CHU de Bordeaux, Place Amélie Raba Léon, 33000 Bordeaux, France. ⁴Institut National d'Etudes Démographiques (INED), Aubervilliers, France. ⁵Inserm, Paris, France. ⁶Etablissement Français du Sang (EFS), Paris, France. ⁷Santé Publique France, French National Public Health Agency, 94415 Saint-Maurice, France. ⁸Research Unit on Children's Psychosocial Maladjustment, Montreal, QC, Canada. ⁹Inserm, UVSQ, CESP, Fac. de Médecine – Université Paris-Sud, INSERM 1018, Paris-Saclay University, DevPsy, Villejuif, France. ¹⁰Public Health and Epidemiology Department, AP-HP Paris-Saclay, Bicêtre Hospital, Le Kremlin-Bicêtre, France. ¹¹Université Paris Cité, Paris, France. ¹²Centre for Research in Epidemiology and Statistics (CRESS), Paris, France. ¹³Institut national de recherche pour l'agriculture, l'alimentation et l'environnement (INRAE), Paris, France. ¹⁴Faculté de Médecine St Antoine, Institut Pierre Louis d'Epidémiologie et de Santé Publique (IPLESP), Equipe de Recherche en Epidémiologie Sociale (ERES), Sorbonne Université, Paris, France. *Lists of authors and their affiliations appear at the end of the paper. email: maeva.monnier@outlook.com; cedric.galera@u-bordeaux.fr

parents have symptoms of anxiety and/or depression have an increased likelihood of symptoms of hyperactivity/inattention and emotional symptoms during school closures in France due to COVID-19. Our findings suggest that public health initiatives should target parents and children to limit the impact of such crises on their mental health issues.

During the pandemic of CoronaVirus Disease first appeared in 2019 (COVID-19), family life suddenly changed with the need to adapt to the restrictive collective measures implemented to prevent the spread of the virus¹. This unexpected situation fraught with uncertainty significantly increased the risk of mental health issues both in parents and children^{1–9}. Several risk factors for parental mental health issues have been identified, such as having a pre-existing mental health condition, parenting a child with special needs, unemployment, single parenting, gender, and financial difficulties^{3,10–12}. COVID-19-related school closures increased the burden on the family, generating a higher risk of parenting-related exhaustion and disrupted parent–child relationships^{8,13,14}. In this stressful situation, the mental health issues of parents and children interacted more than ever^{4,8}.

Upstream of the COVID-19 context, several studies found an increased risk of psychiatric symptoms in parents of children with psychiatric disorders themselves^{15–20}. Some found the same symptoms in parents and children^{15,18}. A Dutch study of 1866 children and their parents revealed that parental depressive and Attention Deficit Hyperactivity Disorder (ADHD) symptoms were further predicted by offspring depression and offspring ADHD, respectively¹⁵. Another study found that parents' symptoms were not always equivalent to their children's mental health issues¹⁷. An Italian case–control study of the parents of 50 children affected by ADHD and of 45 age- and gender-matched healthy children investigated parental psychiatric disorders in ADHD children¹⁷. Compared to parents of children without mental health issues, parents of ADHD children reported higher levels of depression (Odds ratio (OR) 2.36; 95% Confidence Interval (CI) 1.31–4.26) and anxiety disorder (OR 2.01; 95% CI 1.05–3.79)¹⁷. An American retrospective study of 3- to 7-year-old children with ADHD (n = 98) and non-ADHD children (n = 116) showed a greater prevalence of depression and anxiety in parents of children with ADHD²⁰. The same study found that ADHD in the children was associated with higher odds of both maternal and paternal childhood ADHD²⁰.

Furthermore, several studies have identified parental mental health issues (e.g. symptoms of anxiety, depression, ADHD) as a predictor of child mental health issues^{20–23}. A prospective cohort study identified parental mental health issues as one of the strongest predictors of major depressive disorder in 2519 college students²¹. A recent study of 8002 children aged 9–11 years living in the United States found that after adjustment on age, gender, ethnicity, and environmental factors, greater parent mental health issues (i.e., anxious/depressed, withdrawn, somatic complaints, thought problems, attention problems, aggressive behavior, rule-breaking behavior, and intrusive) was associated with greater child mental health issues (i.e., aggressive behavior, anxiety/depression, attention problems, rule-breaking behavior, somatic complaints, social problems, thought problems, and withdrawn/depressed episodes)²².

Although many studies have shown strong associations between parental and child mental health issues, few studies have investigated this relationship during the COVID-19 pandemic, a period of unprecedented impact on the mental health of parents and children^{5,6,24–27}. A cross-sectional survey of 1469 parents in the United Arab Emirates found that parents with severe anxiety levels were seven times more likely to report emotional problems in their children²⁸. A large-scale cross-sectional population study of 29,202 Hong Kong families with children aged 2–12 years explored child psychosocial wellbeing, parent–child interactions, and parental stress during school closures due to COVID-19²⁹. It showed that parents' mental health issues was positively associated with children's hyperactivity/inattention and children's emotional disorder²⁹. An Italian survey of 463 parents found that parental stress (assessed with the Parent Stress Scale) was a risk factor of children's well-being (assessed with the Strengths and Difficulties Questionnaire) during the COVID-19 quarantine³⁰. A French cohort study of 432 families found that after adjustment on child's sex, age, sleeping difficulties, financial difficulties, job situation during lockdown, parents' symptoms of anxiety-depression were associated with children's emotional difficulties (OR 5.7; 95% CI 2.4–5.1) and with children's symptoms of hyperactivity/inattention (OR 2.5; 95% CI 1.4–4.2) during the COVID-19 lockdown³.

Our first objective was to estimate the prevalence of anxiety and depressive symptoms in parents as well as hyperactivity/inattention and emotional symptoms in children during the COVID-19 pandemic. Our second objective was to study the association between parents' mental health issues and children's during the COVID-19 school closure in France, that might have been highlighted during the pandemic. We hypothesized an increased risk of abnormal hyperactivity/inattention and emotional symptoms in children of parents with a high level of anxiety and/or depression symptoms.

Methods

This research is based on the SAPRIS (“SAnité, Perception, pratiques, Relations et Inégalités Sociales en population générale pendant la crise COVID-19”) study. SAPRIS was set up to study the main epidemiological, social and behavioral challenges of the COVID-19 pandemic in France in relation to social inequalities in health and healthcare. More details on this project are available elsewhere³¹. SAPRIS is based on a questionnaire sent to participants of five large epidemiological French cohorts. For this research, we used data pertaining to 3496 children aged 8–9 years old in 2020 and participating in ELFE (Etude Longitudinale Française depuis l'Enfance, The French Longitudinal Study since Childhood) population-based birth cohort that focuses on the health of children born in 2011. All parents in the ELFE cohort not lost at follow-up in April 1, 2020, were contacted³¹. The SAPRIS survey does not have any additional inclusion criteria beyond those original to the ELFE cohort (i.e., give birth in one of the 320 French maternal unit, single or twin live births at 33 weeks of gestation, mother

18 years old, no plan to leave metropolitan France within 3 years and informed consent signed)³². The entire questionnaire is completed by one of the parents between April 16 and June 21, 2020, during the period of school closure in France due to COVID-19. In this study, 3496 children with complete data on their own and their parents' mental health were included in this study (see Fig. 1). All methods were carried out in accordance with the relevant guidelines and regulations. Ethical approval and written informed consents were obtained from all subjects and/or their legal guardian. The Inserm ethics evaluation committee (no 2020.04.24 bis_20.04.22.74247, 2020 April 27), and the CNIL (no 920193, 2020 April 30) approved the research.

Main outcome: children's mental health issues. Two subscales of the French version of the Strengths and Difficulties Questionnaire (SDQ) were used to assess children's scores of hyperactivity/inattention and emotional symptoms^{33,34}. Five items were used to assess each subscale. Parents were asked to rate each of the items regarding their child for the past 15 days on a three-point Likert scale (0. not true to 2. certainly true). From the five items on each subscale, a global score was calculated from 0 to 10. Regarding hyperactivity/inattention subscale, a score ≤ 5 is considered normal, equal to 6 as borderline, and > 6 as abnormal. For emotional symptoms, a score ≤ 3 is considered normal, equal to 4 as borderline, and > 4 as abnormal.

Main exposure: parents' mental health issues. Two measures of parental mental health issues were collected. The first assessed symptoms of anxiety through the Generalized Anxiety Disorder-7 (GAD-7)^{35,36}. This is a 7-item scale scored 0 to 3 (0. never, 1. several days, 2. more than half the time or 3. almost every day), providing a 0 to 21 severity score. Responses were classified as follows: absence of anxiety (< 4), mild anxiety (≤ 9), moderate to severe anxiety (> 9).

A second measure assessed depressive symptom severity with the Patient Health Questionnaire-9 (PHQ-9)^{37,38}. This module includes 9 items on which the subject estimates the frequency of depressive symptomatology (0. never, 1. several days, 2. more than half the time or 3. almost every day). The total PHQ-9 score varies between 0 and 27. Levels of depression are categorized as follows: minimal (score < 5), mild (< 10), moderate to severe (≥ 10)³⁹.

Covariates. Several covariates based on the literature review were taken into account in the statistical analyses, including the child's sleeping difficulties (yes vs no), child's sex (male vs female), sex of the responding parent (male vs female), if the child lived with both parents (yes vs no), parents' occupational grade (executive, intermediate and executive, intermediate and employee, independent, laborer, both inactive or only one employee/laborer), household income per month before lockdown in euros (< 2000 , $[2000; 4000[$, ≥ 4000), and perceived financial situation since the lockdown (stable vs declining).




Figure 1. Flowchart of ELFE/SAPRIS cohort during school closure.

Statistical analysis. Chi-square tests were used to compare the characteristics of the ELFE/SAPRIS cohort children included vs not included in this study. Descriptive statistics (percentages) were used to describe the sample demographic characteristics. Classification and Regression Tree methods (CART) were used to perform multiple imputation to handle missing data⁴⁰. CART methods are useful for imputation because they are robust against outliers, can be applied to mixed data (both continuous and categorical), and can deal with multicollinearity^{40,41}. To test whether parents' mental health influenced their children's mental health, we conducted unadjusted and adjusted multinomial logistic regressions, with hyperactivity/inattention symptoms or emotional symptoms as the dependent variable and depressed or anxious status of parents as predictors. Multinomial logistic regression models were adjusted on the child's emotional symptoms or hyperactivity/inattention symptoms, sleeping difficulties sex, on the respondent's sex, if the child lived with both parents, the parents' occupational category, household income per month before lockdown and perceived financial situation since the lockdown.

Data were analyzed using the *nnet* package in *R* (version 3.6.1) with multinomial logistic regressions specified using the *multinom* function. The *mice* package and its *cart* method were used to perform multiple imputation. The following variables were imputed: child's sex (0.77% of NA), household income per month before lockdown (24.63% of NA), perceived financial situation since the lockdown (4.52% of NA), child's sleeping difficulties (0.11% of NA), and parents' occupational category (0.83% of NA).

Results

Table 1 shows the characteristics of the ELFE/SAPRIS cohort children who were included or not in this study. The two populations did not differ significantly on most variables. They differed only in the number of children living with both parents and in the parents' occupational category. There was a lower proportion of children living with both parents as well as children of parents of high occupational grade. The association between the mental health issues of parents and children might even more pronounced in the population of non-included children.

Table 2 shows the characteristics of the 3496 children included in the study. Of these children, 50% are girls (1736) and 89% (3113) lived with both parents during lockdown. 11.8% (411) of the children had an abnormal hyperactivity/inattention score and 6.6% (231) had an abnormal emotional symptoms score during the school closures.

Most parents (91.9%) who participated were mothers ($n = 3211$). Among responding parents, 7.1% (247) had moderate to severe anxiety, and 6.7% (234) had moderate to severe depression (Table 2).

Parent's mental health issues and children's hyperactivity/inattention. Table 3 shows the adjusted and unadjusted associations between parents' mental health issues and children's symptoms of hyperactivity/inattention. Having parents with moderate to severe anxiety or mild anxiety during school closures was associated with an increased risk of an abnormal hyperactivity/inattention score in children (OR 4.89; 95% CI 3.53–6.76 and OR 2.87; 95% CI 2.27–3.61, respectively) as well as a borderline score (OR 2.51; 95% CI 1.66–3.78 and OR 1.94; 95% CI 1.49–2.52, respectively). These associations remained statistically significant in the multivariate regression model. Adjusted on covariates, moderate to severe anxiety and mild anxiety in the responding parent during school closures were associated with an abnormal hyperactivity-inattention score (aOR 3.31; 95% CI 2.33–4.70 and aOR 2.25; 95% CI 1.76–2.88, respectively) and its borderline level (aOR 2.18; 95% CI 1.42–3.34 and aOR 1.81; 95% CI 1.38–2.37 respectively).

Furthermore, having parents with moderate to severe depression or mild depression during school closure was associated with an increased likelihood of abnormal hyperactivity/inattention scores (OR 6.99; 95% CI 5.07–9.65 & OR 3.21; 95% CI 2.54–4.05, respectively) in children; the same was observed for borderline scores (OR 2.29; 95% CI 1.44–3.64 & OR 2.48; 95% CI 1.93–3.19, respectively) (Table 3). These results remained significant in the multivariate analysis. Adjusting on covariates, moderate to severe parental depression and mild depression during school closures were associated with abnormal hyperactivity-inattention scores (aOR 4.65; 95% CI 3.27–6.59 & aOR 2.54; 95% CI 1.99–3.25, respectively) and borderline scores in children (aOR 2.00; 95% CI 1.24–3.23 & aOR 2.26; 95% CI 1.74–2.93, respectively).

Parents' mental health issues and children's emotional symptoms. Table 4 shows the adjusted and unadjusted associations between parents' mental health issues and children's emotional symptoms scores. Having a parent with moderate to severe anxiety or mild anxiety during school closure was associated with an increased risk of an abnormal emotional symptoms score in children (OR 5.81; 95% CI 3.98–8.61 and OR 3.50; 95% CI 2.59–4.72, respectively) as well as its borderline score (OR 3.11; 95% CI 1.95–4.95 and OR 2.38; 95% CI 1.73–3.27, respectively). Adjusting on covariates, moderate to severe anxiety and mild anxiety in the responding parent during school closures remained associated with an abnormal emotional symptoms score (aOR 3.58; 95% CI 2.33–5.49 and aOR 2.52; 95% CI 1.84–3.46, respectively) and its borderline score (aOR 2.12; 95% CI 1.30–3.46 and aOR 1.98; 95% CI 1.43–2.75, respectively).

Having parents with mild to severe depression during school closures was associated with an increased risk of an abnormal emotional symptoms score in children (OR 6.69; 95% CI 4.56–9.80 and OR 2.77; 95% CI 2.05–3.74, respectively) as well as its borderline score (OR 2.62; 95% CI 1.58–4.34 and OR 2.22; 95% CI 1.63–3.03, respectively) (Table 4). Adjusting on covariates, moderate to severe depression and mild depression in the responding parent during school closures remained associated with an abnormal emotional symptoms score (aOR 5.84; 95% CI 3.91–8.72 and aOR 2.70; 95% CI 1.98–3.68, respectively) and its borderline score (aOR 2.47; 95% CI 1.47–4.15 and aOR 2.28; 95% CI 1.66–3.13, respectively).

	Children included (n = 3496)	Children excluded (n = 1408)	p-value
	n (%)	n (%)	
Child's sex	3469 (99.2)	1395 (99.1)	0.93
Female	1736 (50.0)	700 (50.2)	
Male	1733 (50.0)	695 (49.8)	
Sex of responding parent	3496 (100.0)	1376 (97.7)	0.26
Female	3211 (91.8)	1277 (92.8%)	
Male	285 (8.2)	99 (7.2%)	
Child's sleeping difficulties	3492 (99.9)	1239 (88.0)	0.39
Yes	1349 (38.6)	496 (40.0)	
No	2143 (61.4)	743 (60.0)	
Child lives with both parents	3496 (100.0)	1387 (98.5%)	< 0.05*
Yes	3113 (89.0)	1122 (80.9)	
No	383 (11.0)	265 (19.1)	
Parents' occupational category	3467 (99.2)	1355 (96.2)	< 0.05*
Executive	1145 (33.0)	388 (28.6)	
Intermediate and executive	953 (27.5)	372 (27.5)	
Intermediate and employee	818 (23.6)	334 (24.6)	
Independent	297 (8.6)	121 (8.9)	
Laborer	183 (5.3)	90 (6.6)	
Both inactive or only one employee/laborer	71 (2.0)	50 (3.7)	
Household income per month before lockdown (in euros)	2635 (75.4)	217 (15.4)	0.66
< 2000	183 (6.9)	15 (6.9)	
[2000; 4000[1150 (43.6)	88 (40.6)	
≥ 4000	1302 (49.4)	114 (52.5)	
Perceived financial situation since lockdown	3338 (95.5)	1217 (86.4)	0.25
Declining	910 (27.3)	353 (29.0)	
Stable or increasing	2428 (72.7)	864 (71.0)	
Child's mental health issues			
Hyperactivity/inattention	3496 (100.0)	1153 (81.9)	0.30
Abnormal	411 (11.8)	150 (13.0)	
Borderline	306 (8.8)	88 (7.6)	
Normal	2779 (79.5)	915 (79.4)	
Emotional symptoms	3496 (100.0)	1171 (83.2)	0.22
Abnormal	231 (6.6)	94 (8.0)	
Borderline	196 (5.6)	60 (5.1)	
Normal	3069 (87.8)	1017 (86.8)	
Mental health issues of responding parent			
Anxiety	3496 (100.0)	214 (15.2)	0.55
Moderate to severe	247 (7.1)	18 (8.4)	
Mild	831 (23.8)	45 (21.0)	
Minimal	2418 (69.2)	151 (70.6)	
Depression	3496 (100.0)	188 (13.4)	0.46
Moderate to severe	234 (6.7)	12 (6.4)	
Mild	919 (26.3)	42 (22.3)	
Minimal	2343 (67.0)	134 (71.3)	

Table 1. Comparison of characteristics of children in cohort SAPRIS/ELFE included vs. excluded in this study (n = 4904).

Discussion

This study shows that moderate to severe anxiety and moderate to severe depression in parents during school closures were associated with both abnormal hyperactivity-inattention and emotional symptoms scores in children. Furthermore, mild anxiety and mild depression in the responding parent during school closures were associated with abnormal scores of both hyperactivity-inattention and emotional symptoms in children.

Although the impact of COVID-19 on the decline of mental health remains debated up to date^{42–44}, our results are consistent with previous research showing that parental mental health issues during the pandemic were associated with children's mental health issues^{3,8,45}. A longitudinal Canadian study of 68 families of children aged

	n (%)
Child's sex	3469 (99.2)
Female	1736 (50.0)
Male	1733 (50.0)
Sex of responding parent	3496 (100.0)
Female	3211 (91.8)
Male	285 (8.2)
Child's sleeping difficulties	3492 (99.9)
Yes	1349 (38.6)
No	2143 (61.4)
Child lives with both parents	3496 (100.0)
Yes	3113 (89.0)
No	383 (11.0)
Parents' occupational category	3467 (99.2)
Executive	1145 (33.0)
Intermediate and executive	953 (27.5)
Intermediate and employee	818 (23.6)
Independent	297 (8.6)
Laborer	183 (5.3)
Both inactive or only one employee/laborer	71 (2.0)
Household income per month before lockdown (in euros)	2635 (75.4)
< 2000	183 (6.9)
[2000; 4000[1150 (43.6)
≥ 4000	1302 (49.4)
Perceived financial situation since lockdown	3338 (95.5)
Declining	910 (27.3)
Stable or increasing	2428 (72.7)
Child's mental health issues	
Hyperactivity/inattention	3496 (100.0)
Abnormal	411 (11.8)
Borderline	306 (8.8)
Normal	2779 (79.5)
Emotional symptoms	3496 (100.0)
Abnormal	231 (6.6)
Borderline	196 (5.6)
Normal	3069 (87.8)
Mental health issues of responding parent	
Anxiety	3496 (100.0)
Moderate to severe	247 (7.1)
Mild	831 (23.8)
Minimal	2418 (69.2)
Depression	3496 (100.0)
Moderate to severe	234 (6.7)
Mild	919 (26.3)
Minimal	2343 (67.0)

Table 2. Characteristics of children and their families in the ELFE cohort of the SAPRIS study during school closure in France (n = 3496).

7 to 9 years showed that higher levels of maternal anxiety during the pandemic were associated with a greater increase in internalizing problems in children⁸. A cross-sectional Italian study of 721 parents found that parents' behaviors during lockdown (i.e., employment status) were significantly related to their own internalizing symptoms. These symptoms, in turn, had a strong and positive predictive effect on children's internalizing problems⁴⁵.

The cross-sectional associations can be explained by the reciprocal influences of parental and child mental health issues. This study shows that the associations between parents' and children's mental health issues which have already been highlighted outside the COVID context¹⁵⁻²⁰ is found to be also present during the COVID-19 pandemic. Based on the results of this study, we are not able to know whether this association is stronger during the COVID-19 period due to the lack of data prior to the pandemic. However, the lockdown and school closures have impacted families' daily life and have challenged their parenting skills. Dealing with school closure and

	OR		aOR ²	
	Abnormal ¹	Borderline ¹	Abnormal ¹	Borderline ¹
Anxiety of responding parent				
Moderate to severe	4.89 [3.53; 6.76]	2.51 [1.66; 3.78]	3.31 [2.33; 4.70]	2.18 [1.42; 3.34]
Mild	2.87 [2.27; 3.61]	1.94 [1.49; 2.52]	2.25 [1.76; 2.88]	1.81 [1.38; 2.37]
Minimal	–	–	–	–
Depression of responding parent				
Moderate to severe	6.99 [5.07; 9.65]	2.29 [1.44; 3.64]	4.65 [3.27; 6.59]	2.00 [1.24; 3.23]
Mild	3.21 [2.54; 4.05]	2.48 [1.93; 3.19]	2.54 [1.99; 3.25]	2.26 [1.74; 2.93]
Minimal	–	–	–	–

Table 3. Association between parents' mental health issues and children's hyperactivity/inattention score: multinomial logistic regressions (n = 3496). ¹Normal is reference group. ²Results adjusted on child's emotional symptoms score, child's sleeping difficulties, child's sex, sex of responding parent, if the child lived with both parents, parents' occupational category, household income per month before lockdown and perceived financial situation since lockdown.

	OR		aOR ²	
	Abnormal ¹	Borderline ¹	Abnormal ¹	Borderline ¹
Anxiety of responding parent				
Moderate to severe	5.81 [3.98; 8.61]	3.11 [1.95; 4.95]	3.58 [2.33; 5.49]	2.12 [1.30; 3.46]
Mild	3.50 [2.59; 4.72]	2.38 [1.73; 3.27]	2.52 [1.84; 3.46]	1.98 [1.43; 2.75]
Minimal	–	–	–	–
Depression of responding parent				
Moderate to severe	6.69 [4.56; 9.80]	2.62 [1.58; 4.34]	3.78 [2.47; 5.78]	1.79 [1.05; 3.05]
Mild	2.77 [2.05; 3.74]	2.22 [1.63; 3.03]	1.98 [1.44; 2.73]	1.78 [1.29; 2.46]
Minimal	–	–	–	–

Table 4. Association between parents' mental health issues and children's emotional symptoms score: multinomial logistic regressions (n = 3496). ¹Normal score is reference group. ²Results adjusted on child's hyperactivity/inattention score, child's sleeping difficulties, child's sex, sex of responding parent, if the child lived with both parents, parents' occupational category, household income per month before lockdown and perceived financial situation since lockdown.

the lockdown is a particularly stressful situation for parents who have to balance personal life, work, and ensure the continuity of children's education and learning¹. This situation puts parents at a higher risk of experiencing anxiety, stress, depression, potentially impairing their ability to be supportive caregivers¹. We speculate that more anxiety and depression in parents prevents them from understanding their child's needs and responding appropriately. Moreover, the COVID-19 pandemic was marked by an increase in irritability, shouting at children, verbal abuse, and parents punishing children^{46–48}. Such behaviors are associated with an increased risk of disorders in children^{49–51}. On the other hand, managing children with hyperactivity/inattention and/or emotional symptoms can be a source of stress and mental distress for parents^{14,52}. An Italian study of 1226 parents showed that having a child with cognitive or physical disabilities was a risk factor of parenting-related exhaustion during COVID-19 lockdown¹⁴. Furthermore, a recent cohort study of 917 parents showed that more child emotional and hyperactivity-inattention symptoms were significant predictors of parental distress during the lockdown⁵². Despite these environmental factors that may explain the parents' and children's mental health issues mutually influencing each other, the hypothesis of a genetic vulnerability cannot be ruled out. Indeed, several studies produced evidence for genetic overlap between ADHD and bipolar disorder^{53,54}, and ADHD and major depressive disorder^{53–55}. Studies taking into account both environmental risk factors and genes are needed to better understand the relationships between various mental health issues in parents and offspring.

Our research has several strengths. First, the sample was large with sufficient power, thus increasing the likelihood of obtaining more conclusive results with adjustments. In addition, the measures of parental and children's mental health were based on validated scales with good psychometric qualities. Moreover, we took several confounding factors into account such as the sex of the child and the responding parent, the occupational category of the household and the perceived financial situation.

However, several limitations must be taken into consideration. First, we were not able to assess a potential effect of school closures due to the absence of repeated measures of children's and parents' mental health issues before the lockdown. Due to the lack of measurement of parents' and children's mental health issues just prior to the pandemic, we are unable to identify an effect of the pandemic on the parents' and children's mental health issues association. On the other hand, this study focuses on children aged 8–9 years; other studies are needed on

other age groups to be able to generalize the results to other age ranges. In addition, we were unable to infer any causal relationship between parents' and children's mental health issues. Also, the sample was composed mostly of mothers. However, we adjusted the analyses for the gender of the responding parent to make the results generalizable. Furthermore, we didn't have a measure of parental stress, which is an essential variable when studying parents' mental health issues, especially during the period of COVID-19³⁰. At last, these are only self-reported measures and thus are subject to bias in the assessment. To avoid the later, children's mental health issues should be measured by a health care professional. Parents who are anxious or depressed may be more likely to indicate a mental disorder in their child than healthy parents.

For future research, we recommend to have repeated measures and study a wider range of mental health issues in parents (e.g., mood disorders, substance use disorders, and psychotic disorders) and children (e.g., learning disabilities, autism spectrum disorders). It could also be interesting to study how the association between parents' and children's mental health issues evolves once the school takes over.

The present findings have significant implications. Parenting while having a mental health issues during school closure may represent a threat to children's mental health issues and have an impact on the way children and parents interact. Governments and healthcare authorities should be alerted to the effects that lockdown and school closure may worsen symptoms in children and their parents. Whenever mental health issues are suspected in the child or their parents, it is of utmost importance that clinicians take into consideration the mental health of the other family members close to the patient. In the same line, prevention strategies promoting parental mental health issues could have the potential for positive long-term impacts on child mental health issues.

Longitudinal studies are now needed to assess changes in the mental health of children and parents during and after the pandemic. These results underline the importance of focusing on mental health issues in children in the presence of parental mental health issues and vice-versa. This seems to be even truer in this period of collective stress and promiscuity characteristic of school closures and the lockdown in France due to the COVID-19 pandemic. In addition, continuing to deliver care during the crisis has posed problems. One of our previous studies showed that 87% of children receiving psychological care prior to the pandemic were unable to continue to receive it⁵⁶. More thorough follow-up of children and parents at risk is required to ensure continuity of care during such periods of risk. Finally, the creation of a national cohort would allow retrospective and prospective data to be gathered to study the evolution of mental health in France over time and during crises such as the COVID-19 pandemic.

Conclusion

Children of parents with anxiety and/or depression were at increased risk of hyperactivity/inattention and emotional symptoms during school closures in France due to COVID-19. The COVID-19 pandemic and associated school closures have caused disruption to family life and affected the wellbeing of children and their parents. Findings suggest that public health initiatives should target parents and children to limit the impact of such crises on their mental health issues.

Data availability

The study data are protected by the health data privacy regulations set out by the French National Commission on Informatics and Liberty (Commission Nationale de l'Informatique et des Libertés, CNIL). The data are available upon reasonable request after consultation with the Sapis steering committee. While French law forbids free access to the Sapis data, it may be granted by the steering committee after legal verification of how the data will be used. The datasets generated during and/or analysed during the current study are available from Dr Cédric Galéra on reasonable request.

Received: 13 October 2022; Accepted: 21 March 2023

Published online: 24 March 2023

References

1. Spinelli, M., Lionetti, F., Pastore, M. & Fasolo, M. Parents' stress and children's psychological problems in families facing the COVID-19 Outbreak in Italy. *Front. Psychol.* **11**, 1713 (2020).
2. Nearchou, F., Flinn, C., Niland, R., Subramaniam, S. S. & Hennessy, E. Exploring the impact of COVID-19 on mental health outcomes in children and adolescents: A systematic review. *Int. J. Environ. Res. Public Health* **17**(22), E8479 (2020).
3. Moulin, F. *et al.* Risk and protective factors related to children's symptoms of emotional difficulties and hyperactivity/inattention during the COVID-19-related lockdown in France: Results from a community sample. *Eur. Child Adolesc. Psychiatry* **31**, 1–12 (2021).
4. Kim, S. J. *et al.* Parental mental health and children's behaviors and media usage during COVID-19-related school closures. *J. Korean Med. Sci.* **36**(25), e184 (2021).
5. Jacques-Aviñó, C. *et al.* Gender-based approach on the social impact and mental health in Spain during COVID-19 lockdown: A cross-sectional study. *BMJ Open* **10**(11), e044617 (2020).
6. Fiorillo, A. *et al.* Effects of the lockdown on the mental health of the general population during the COVID-19 pandemic in Italy: Results from the COMET collaborative network. *Eur. Psychiatry J. Assoc. Eur. Psychiatry* **63**(1), e87 (2020).
7. Yamamura, E. & Tsutsui, Y. School closures and mental health during the COVID-19 pandemic in Japan. *J. Popul. Econ.* **19**, 1–38 (2021).
8. Khoury, J. E., Kaur, H. & Gonzalez, A. Parental mental health and hostility are associated with longitudinal increases in child internalizing and externalizing problems during COVID-19. *Front. Psychol.* **12**, 706168 (2021).
9. Ramiz, L. *et al.* A longitudinal study of mental health before and during COVID-19 lockdown in the French population. *Glob. Health* **17**(1), 29 (2021).
10. Amorim, R. *et al.* The impact of COVID-19 on children with autism spectrum disorder. *Rev. Neurol.* **71**(8), 285–291 (2020).
11. El-Osta, A. *et al.* How is the COVID-19 lockdown impacting the mental health of parents of school-age children in the UK? A cross-sectional online survey. *BMJ Open* **11**(5), e043397 (2021).

12. Cao, W. *et al.* The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry Res.* **287**, 112934 (2020).
13. Calvano, C. *et al.* Families in the COVID-19 pandemic: Parental stress, parent mental health and the occurrence of adverse childhood experiences—Results of a representative survey in Germany. *Eur. Child Adolesc. Psychiatry* **31**, 1–13 (2021).
14. Marchetti, D. *et al.* Parenting-related exhaustion during the Italian COVID-19 lockdown. *J. Pediatr. Psychol.* **45**(10), 1114–1123 (2020).
15. Wesseldijk, L. W. *et al.* Risk factors for parental psychopathology: A study in families with children or adolescents with psychopathology. *Eur. Child Adolesc. Psychiatry* **27**(12), 1575–1584 (2018).
16. Vidair, H. B. *et al.* Screening parents during child evaluations: Exploring parent and child psychopathology in the same clinic. *J. Am. Acad. Child Adolesc. Psychiatry* **50**(5), 441–450 (2011).
17. Margari, F. *et al.* Parents psychopathology of children with attention deficit hyperactivity disorder. *Res. Dev. Disabil.* **34**(3), 1036–1043 (2013).
18. Segenreich, D. *et al.* Multilevel analysis of ADHD, anxiety and depression symptoms aggregation in families. *Eur. Child Adolesc. Psychiatry* **24**(5), 525–536 (2015).
19. Gao, Y. *et al.* Resemblance and clustering of mother's and father's psychopathology levels among Chinese parents of schoolchildren with psychiatric disorders. *Eur. Psychiatry J. Assoc. Eur. Psychiatry* **63**(1), e100 (2020).
20. Chronis, A. M. *et al.* Psychopathology and substance abuse in parents of young children with attention-deficit/hyperactivity disorder. *J. Am. Acad. Child Adolesc. Psychiatry* **42**(12), 1424–1432 (2003).
21. Ebert, D. D. *et al.* Prediction of major depressive disorder onset in college students. *Depress Anxiety* **36**(4), 294–304 (2019).
22. Zhang, H., Lee, Z. X., White, T. & Qiu, A. Parental and social factors in relation to child psychopathology, behavior, and cognitive function. *Transl. Psychiatry* **10**(1), 80 (2020).
23. Breaux, R. P., Harvey, E. A. & Lugo-Candelas, C. I. The role of parent psychopathology in the development of preschool children with behavior problems. *J. Clin. Child Adolesc. Psychol.* **43**(5), 777–790 (2014).
24. Gloster, A. T. *et al.* Impact of COVID-19 pandemic on mental health: An international study. *PLoS ONE* **15**(12), e0244809 (2020).
25. Alkhamees, A. A., Alrashed, S. A., Alzunaydi, A. A., Almohimeed, A. S. & Aljohani, M. S. The psychological impact of COVID-19 pandemic on the general population of Saudi Arabia. *Compr. Psychiatry* **102**, 152192 (2020).
26. Wang, C. *et al.* Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *Int. J. Environ. Res. Public Health* **17**(5), E1729 (2020).
27. Tee, M. L. *et al.* Psychological impact of COVID-19 pandemic in the Philippines. *J. Affect. Disord.* **277**, 379–391 (2020).
28. Saddik, B. *et al.* The psychological impact of the COVID-19 pandemic on adults and children in the United Arab Emirates: A nationwide cross-sectional study. *BMC Psychiatry* **21**, 224 (2021).
29. Tso, W. W. Y. *et al.* Vulnerability and resilience in children during the COVID-19 pandemic. *Eur. Child Adolesc. Psychiatry* **31**(1), 161–176 (2020).
30. Cusinato, M. *et al.* Stress, resilience, and well-being in Italian children and their parents during the COVID-19 pandemic. *Int. J. Environ. Res. Public Health* **17**(22), E8297 (2020).
31. Carrat, F. *et al.* Incidence and risk factors of COVID-19-like symptoms in the French general population during the lockdown period: A multi-cohort study. *BMC Infect. Dis.* **21**, 169 (2021).
32. Charles, M.-A. *et al.* Cohort Profile: The French National cohort of children ELFE: Birth to 5 years. *Int. J. Epidemiol.* **49**(2), 368–369 (2020).
33. Goodman, R. Psychometric properties of the strengths and difficulties questionnaire. *J. Am. Acad. Child Adolesc. Psychiatry* **40**(11), 1337–1345 (2001).
34. Shojaei, T., Wazana, A., Pitrou, I. & Kovess, V. The strengths and difficulties questionnaire: Validation study in French school-aged children and cross-cultural comparisons. *Soc. Psychiatry Psychiatr. Epidemiol.* **44**(9), 740–747 (2009).
35. Spitzer, R. L., Kroenke, K., Williams, J. B. W. & Löwe, B. A brief measure for assessing generalized anxiety disorder: The GAD-7. *Arch. Intern. Med.* **166**(10), 1092–1097 (2006).
36. Spitzer, R. L., Williams, J. & Kroenke, K. *French Version of the GAD-7*. <https://medfam.umontreal.ca/wp-content/uploads/sites/16/GAD-7-fran%C3%A7ais.pdf>.
37. Spitzer, R. L., Kroenke, K. & Williams, J. B. Validation and utility of a self-report version of PRIME-MD: The PHQ primary care study. Primary care evaluation of mental disorders. Patient health questionnaire. *JAMA* **282**(18), 1737–44 (1999).
38. Carballeira, Y. *et al.* Criterion validity of the French version of patient health questionnaire (PHQ) in a hospital department of internal medicine. *Psychol. Psychother.* **80**(1), 69–77 (2007).
39. Kroenke, K., Spitzer, R. L. & Williams, J. B. The PHQ-9: Validity of a brief depression severity measure. *J. Gen. Intern. Med.* **16**(9), 606–613 (2001).
40. Burgette, L. F. & Reiter, J. P. Multiple imputation for missing data via sequential regression trees. *Am. J. Epidemiol.* **172**(9), 1070–1076 (2010).
41. Van Buuren, S. *Flexible Imputation of Missing Data* 2nd edn. (Chapman & Hall, 2012).
42. Penninx, B. W. J. H. *et al.* How COVID-19 shaped mental health: From infection to pandemic effects. *Nat. Med.* **28**, 2027–2037 (2022).
43. Wu, T. *et al.* Prevalence of mental health problems during the COVID-19 pandemic: A systematic review and meta-analysis. *J. Affect. Disord.* **15**(281), 91–98 (2021).
44. Robinson, E., Sutin, A. R., Daly, M. & Jones, A. A systematic review and meta-analysis of longitudinal cohort studies comparing mental health before versus during the COVID-19 pandemic in 2020. *J. Affect. Disord.* **296**, 567–576 (2022).
45. Crescentini, C. *et al.* Stuck outside and inside: An exploratory study on the effects of the COVID-19 outbreak on Italian parents and children's internalizing symptoms. *Front. Psychol.* **11**, 586074 (2020).
46. Shah, R., Raju, V. V., Sharma, A. & Grover, S. Impact of COVID-19 and lockdown on children with ADHD and their families—An online survey and a continuity care model. *J. Neurosci. Rural Pract.* **12**(1), 71–79 (2021).
47. Lee, S. J., Ward, K. P., Lee, J. Y. & Rodriguez, C. M. Parental social isolation and child maltreatment risk during the COVID-19 pandemic. *J. Fam. Violence* **37**(5), 813–824 (2022).
48. Norman, R. E. *et al.* The long-term health consequences of child physical abuse, emotional abuse, and neglect: A systematic review and meta-analysis. *PLoS Med.* **9**(11), e1001349 (2012).
49. Beckmann, L. Does parental warmth buffer the relationship between parent-to-child physical and verbal aggression and adolescent behavioural and emotional adjustment?. *J. Fam. Stud.* **27**(3), 366–387 (2021).
50. Crum, K. I. & Moreland, A. D. Parental stress and children's social and behavioral outcomes: The role of abuse potential over time. *J. Child Fam. Stud.* **26**(11), 3067–3078 (2017).
51. Font, S. A. & Cage, J. Dimensions of physical punishment and their associations with children's cognitive performance and school adjustment. *Child Abuse Negl.* **75**, 29–40 (2018).
52. Mazza, C. *et al.* The COVID-19 lockdown and psychological distress among Italian parents: Influence of parental role, parent personality, and child difficulties. *Int. J. Psychol. J. Int. Psychol.* **56**(4), 577–584 (2021).
53. Hammerschlag, A. R., De Leeuw, C. A., Middeldorp, C. M. & Polderman, T. J. C. Synaptic and brain-expressed gene sets relate to the shared genetic risk across five psychiatric disorders. *Psychol. Med.* **50**(10), 1695–1705 (2020).

54. Zhao, H. & Nyholt, D. R. Gene-based analyses reveal novel genetic overlap and allelic heterogeneity across five major psychiatric disorders. *Hum. Genet.* **136**(2), 263–274 (2017).
55. Cross-Disorder Group of the Psychiatric Genomics Consortium, International Inflammatory Bowel Disease Genetics Consortium (IBDGC). Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. *Nat. Genet.* **45**(9), 984–94 (2013).
56. Monnier, M. *et al.* Children's mental and behavioral health, schooling, and socioeconomic characteristics during school closure in France due to COVID-19: The SAPRIS project. *Sci. Rep.* **11**(1), 22373 (2021).

Acknowledgements

The authors are grateful to the Nouvelle-Aquitaine Regional Government and its *AMI FLASH Recherche et Innovations COVID* program, without which the project would never have started. The authors warmly thank all the volunteers of the ELFE cohort. They thank the staff of the SAPRIS study group who worked with dedication and commitment to collect and manage the data and to ensure continuing communication with the cohort participants.

Author contributions

M.M. contributed to the request for data access, literature search, data-management, data analysis, and wrote the first draft of the manuscript. C.G. and M.M. contributed to the data interpretation and writing. All authors reviewed the manuscript. C.G. was the senior author of the manuscript. M.M. and C.G. had access to the data of this study and takes responsibility for the integrity and the accuracy of the data analysis.

Funding

Sponsor and funding sources played no role in the study design, data collection, analysis, interpretation or drafting of the study. MM and CG had full access to all data in the study. ANR (Agence Nationale de la Recherche, 0009/SAPRIS/997/NB), Fondation pour la Recherche Médicale, Direction Générale de la Recherche et de l'Innovation, Institut Gustave Roussy, the Nouvelle-Aquitaine Regional Government.

Competing interests

Mrs Maeva Monnier was funded by the Nouvelle-Aquitaine Regional Government (AMI Flash Recherche et Innovations COVID). The other authors declare no competing interests related to this paper.

Additional information

Correspondence and requests for materials should be addressed to M.M. or C.G.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023

The SAPRIS Study Group

Nathalie Bajos¹⁵, Fabrice Carrat¹⁶, Pierre-Yves Ancel^{10,12,13,14}, Marie-Aline Charles^{4,5,10,11,12,13}, Florence Jusot¹⁷, Claude Martin^{18,19}, Laurence Meyer²⁰, Ariane Pailhé⁴, Alexandra Rouquette^{9,10}, Gianluca Severi^{10,21}, Alexis Spire^{18,19}, Mathilde Touvier^{10,12,13,22} & Marie Zins²³

¹⁵IRIS (UMR 8156 CNRS - EHESS - U997 INSERM), Institut de Recherche Interdisciplinaire sur les Enjeux Sociaux - Sciences sociales, politique, santé, Aubervilliers, France. ¹⁶INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique (IPLESP), Sorbonne Université, Paris, France. ¹⁷Paris-Dauphine University-PSL, Paris, France. ¹⁸French National Centre for Scientific Research (CNRS), Paris, France. ¹⁹National Institute for Demographic Studies, Paris, France. ²⁰University of Paris Sud, Paris, France. ²¹E4N & EN3-Cohorts, Paris, France. ²²Inra U1125, CNAM, Paris 13 University, Nutritional Epidemiology Research Team (EREN), Bobigny, France. ²³Sorbonne University, Paris, France.

The ELFE Joint Unit

Xavier Thierry^{4,5,10,11}, Thierry Simeon^{4,5,10,11}, Bertrand Geay^{4,5,10,11},
Laetitia Marie-Noelle Dufourg^{4,5,10,11} & Marie-Aline Charles^{4,5,10,11,12,13}